

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 111 067 A2**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
27.06.2001 Bulletin 2001/26

(51) Int Cl.7: **C12P 23/00, C12N 9/08,
C12N 9/02**

(21) Application number: **00126114.8**

(22) Date of filing: **29.11.2000**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **01.12.1999 EP 99123821**

(71) Applicant: **F. HOFFMANN-LA ROCHE AG
4070 Basel (CH)**

(72) Inventors:
• **Hoshino, Tatsuo**
Kamakura-shi, Kanagawa-ken 248-0027 (JP)
• **Ojima, Kazuyuki**
Fujisawa-shi, Kanagawa-ken 251-0877 (JP)
• **Setoguchi, Yutaka**
Fujisawa-shi, Kanagawa-ken 251-0033 (JP)

(74) Representative: **Kellenberger, Marcus Dr. et al
Grenzacherstrasse 124
4070 Basel (CH)**

(54) **Recombinant production of carotenoids, particularly astaxanthin**

(57) The present invention is directed to a process for producing carotenoids comprising cultivating a recombinant organism having a gene for one or more active oxygen species-quenching factor(s) that is substantially disrupted with a disruption cassette specific to the

gene, and recovering carotenoids from the culture, as well as to genetic materials useful in the said process, such as a recombinant organism producible of carotenoids, a disruption cassette, a recombinant DNA sequence, a recombinant DNA fragment; and a polynucleotide.

EP 1 111 067 A2

BEST AVAILABLE COPY

Description

[0001] The present invention relates to recombinant production of carotenoids, particularly astaxanthin, and biological materials useful therefor.

[0002] Astaxanthin is known to be distributed in a wide variety of organisms such as animal (e.g. birds such as flamingo and scarlet ibis, and fish such as rainbow trout and salmon), algae and microorganisms. It is also recognized that astaxanthin has a strong antioxidation property against active oxygen as well as most of carotenoids, which is expected to apply to pharmaceutical usage to protect living cells against some diseases such as a cancer. Moreover, from a viewpoint of industrial application, a demand for astaxanthin as a coloring reagent is increasing especially in the industry of farmed fish, such as salmon, because astaxanthin imparts distinctive orange-red coloration to the animals and contributes to consumer appeal in the marketplace.

[0003] *Phaffia rhodozyma* is known as a carotenogenic yeast strain which produces astaxanthin specifically. Different from the other carotenogenic yeast, *Rhodotorula* species, *Phaffia rhodozyma* can ferment some sugars such as D-glucose. This is an important feature from a viewpoint of industrial application. In a recent taxonomic study, a sexual cycle of *P. rhodozyma* was revealed and its teleomorphic state was designated under the name of *Xanthophyllomyces dendrorhous* (W.I. Golubev; *Yeast* 11, 101 - 110, 1995). Some strain improvement studies to obtain hyper producers of astaxanthin from *P. rhodozyma* have been conducted, but such efforts have been restricted to employ the method of conventional mutagenesis and protoplast fusion in this decade. Recently, Wery *et al.* developed a host vector system using *P. rhodozyma* in which a non-replicable plasmid was used to be integrated onto the genome of *P. rhodozyma* at the locus of ribosomal DNA in multicopies (Wery *et al.*, *Gene*, 184, 89-97, 1997). And Verdoes *et al.* reported more improved vectors to obtain a transformant of *P. rhodozyma* as well as its three carotenogenic genes which code for the enzymes that catalyze the reactions from geranylgeranyl pyrophosphate to betacarotene (International patent WO97/23633). The importance of genetic engineering method on the strain improvement study of *P. rhodozyma* will increase in near future to break through the reached productivity by the conventional methods.

[0004] As described above, astaxanthin has an antioxidant property and this feature seems to have an important role *in vivo* for a protection against active oxygen species such as O_2^- , H_2O_2 and OH^\cdot , which are continuously generated in living cells. An *et al.* obtained hyperproducer of astaxanthin from *P. rhodozyma* by selection of antimycin-sensitive strain after a conventional chemical mutagenesis (An, G-H. *et al.*, *Appl. Env. Microbiol.*, 55 (1), 116-124, 1989). Antimycin is known to be an inhibitor of respiratory chain between cytochrome b and C_1 (Lucchini, G. *et al.*, *Mol. Gen. Genet.*, 177, 139-, 1979) and enhanced pigmentation in such an antimycin-sensitive mutants. Active oxygen species produced due to a blockade of the primary respiratory chain at the bc_1 complex stimulated carotenoid formation (An, G-H *et al.*, *Appl. Env. Microbiol.*, 55, 116-124, 1989). Indeed, addition of an O_2^- generator, duroquinone, to the growth medium increased total carotenoid content (main carotenoid is astaxanthin) as well as the relative amounts of xanthophylls present in *P. rhodozyma*, while the active oxygen species-quenching factor mannitol reversed this effect (Schroeder, W. A. *et al.*, *J. Gen. Microbiol.*, 139, 907-912, 1993). These results prompted such authors to speculate an antioxidant property of astaxanthin in *P. rhodozyma*. In fact, astaxanthin production is stimulated in post-exponential growth phase when respiration activity is fully induced. Moreover, the addition of respiratory substrate such as ethanol to the medium enhanced the astaxanthin production in *P. rhodozyma* (Gu, W-L. *et al.*, *J. Ind. Microbiol. Biotechnol.*, 19, 114-117, 1997). Although Schroeder *et al.* tried to determine a relationship of activities of superoxide dismutase (SOD) and catalase which act as native active oxygen species-quenching factors in *P. rhodozyma* to the productivity of astaxanthin by comparing a difference between parent strain and antimycin-sensitive hyperproducer of astaxanthin, direct correlation of *in vitro* activity could not be observed.

[0005] In accordance with the present invention the genes and the enzymes for active oxygen species-quenching factor(s), such as SOD and catalase which are biological materials useful in the improvement of the carotenoid, particularly astaxanthin, production process are provided. This invention involves cloning and determination of the genes which code for mitochondrial and cytoplasmic SODs and catalase. This invention also involves the enzymatic characterization as a result of the disruption of the genes in *P. rhodozyma*. Their disruption effects on the carotenogenesis can be confirmed by the cultivation of such transformants in an appropriate medium under an appropriate cultivation condition. More particularly, the present invention provides a process for producing carotenoids which comprises cultivating a recombinant organism whose gene for one or more active oxygen species-quenching factor(s) is substantially disrupted with the aid of a disruption cassette specific to the said gene, and recovering carotenoids from the culture. The host organism of the said recombinant organism may belong to the kingdom of *Monera*, *Protista* or *Fungi*. More preferable host organism of the said recombinant organism may belong to the genus *Erwinia*, *Rhodobacter*, *Myxococcus*, *Flavobacter*, *Paracoccus*, *Synechococcus*, *Synechocystis*, *Agrobacterium*, *Streptomyces*, *Haematococcus*, *Dunaliella*, *Phaffia*, *Xanthophyllomyces*, *Neurospora*, *Rhodotorula*, *Blakeslea*, or *Phycomyces*. Most preferably, the said host organism is a strain of *P. rhodozyma*.

[0006] The active oxygen species-quenching factor(s) is(are) mitochondrial superoxide dismutase (SOD), cytoplasmic superoxide dismutase (SOD) and/or catalase.

[0007] The present invention also provides a recombinant organism producible of carotenoids, which is characterized in that the gene for at least one active oxygen species-quenching factor is substantially disrupted with the aid of introduction of a disruption cassette specific to the said gene. The active oxygen species-quenching factor(s) to be disrupted is(are) mitochondrial superoxide dismutase (SOD), cytoplasmic superoxide dismutase (SOD) and/or catalase.

[0008] The present invention further provides a disruption cassette which can be used to prepare the said recombinant organism of the present invention for disrupting a gene coding for an active oxygen species-quenching factor effective in an organism which is producible of carotenoids, which comprises a partial nucleotide sequence substantially identical to a part of the DNA sequence coding for an active oxygen species-quenching factor and a selectable marker gene. For the construction of the disruption cassette, the target organism may belong to the kingdom of *Monera*, *Protista* or *Fungi*, more preferably to the genus *Erwinia*, *Rhodobacter*, *Myxococcus*, *Flavobacter*, *Paracoccus*, *Synechococcus*, *Synechocystis*, *Agrobacterium*, *Streptomyces*, *Haematococcus*, *Dunaliella*, *Phaffia*, *Wanophyllum*, *Neurospora*, *Rhodotorula*, *Blakeslea*, or *Phycomyces*. The active oxygen species-quenching factor(s) to be disrupted is(are) mitochondrial superoxide dismutase (SOD), cytoplasmic superoxide dismutase (SOD) and/or catalase.

[0009] As used herein, a polynucleotide or polypeptide sequence (A) is said to be substantially identical to another sequence (B) if sequence A is at least 75% identical, preferably 85% identical, such as at least 95% identical to sequence B.

[0010] Further, the present invention provides a recombinant DNA sequence coding for an active oxygen species-quenching factor effective in an organism producible of carotenoids. The DNA sequence may be originated from an organism belonging to the kingdom of *Monera*, *Protista* or *Fungi*, more preferably to the genus *Erwinia*, *Rhodobacter*, *Myxococcus*, *Flavobacter*, *Paracoccus*, *Synechococcus*, *Synechocystis*, *Agrobacterium*, *Streptomyces*, *Haematococcus*, *Dunaliella*, *Phaffia*, *Xanthophyllum*, *Neurospora*, *Rhodotorula*, *Blakeslea*, or *Phycomyces*. Particularly preferred organism is *P. rhodozyma*. The active oxygen species-quenching factor coded by the recombinant DNA sequence may be mitochondrial superoxide dismutase, cytoplasmic superoxide dismutase and/or catalase.

[0011] A recombinant DNA sequence coding for mitochondrial superoxide dismutase can be identified by SEQ ID NO: 1 or 4, or it may have high homology to the sequence of SEQ ID NO: 1 or 4 enough to recombine with either of the sequences of SEQ ID NOs: 1 and 4. A recombinant DNA sequence coding for cytoplasmic superoxide dismutase can be identified by SEQ ID NO: 2 or 6, or it may have high homology to the sequence of SEQ ID NO: 2 or 6 enough to recombine with either of the sequences of SEQ ID NOs: 2 and 6. Also a recombinant DNA sequence coding for catalase can be identified by SEQ ID NO: 3 or 8, or it may have high homology to the sequence of SEQ ID NO: 3 or 8 enough to recombine with either of the sequences of SEQ ID NOs: 3 and 8.

[0012] In the present invention, any combination of the following hybridization and wash conditions may be used, as appropriate, to identify homologous polynucleotide sequences:

High Stringency Hybridization:

[0013]

6X SSC
0.5% SDS
100 ug/ml denatured salmon sperm DNA
50% formamide
Incubate overnight with gentle rocking at 42°C overnight.

High Stringency Wash:

[0014] 1 wash in 2X SSC, 0.5% SDS at Room Temperature for 15 minutes, followed by another wash in 0.1X SSC, 0.5% SDS at Room Temperature for 15 minutes.

Low Stringency Hybridization:

[0015]

6X SSC
0.5% SDS
100 ug/ml denatured salmon sperm DNA
50% formamide
Incubate overnight with gentle rocking at 37°C overnight.

Low Stringency Wash:

[0016] 1 wash in 0.1X SSC, 0.5% SDS at Room Temperature for 15 minutes.

[0017] Moderately stringent conditions may be obtained by varying the temperature at which the hybridization reaction occurs and/or the wash conditions as set forth above.

[0018] Thus, as used herein, a sequence (A) is said to have "high homology" to another sequence (B) if sequence A hybridizes to sequence B under high stringency conditions (*i.e.*, high stringency hybridization and wash conditions as defined above), and if the polypeptide or polypeptide fragment encoded by sequence A has the same activity as the polypeptide encoded by B.

[0019] The present invention further provides a recombinant DNA fragment which comprises a coding region for a transit peptide upstream the coding region of an objective protein which may be mitochondrial superoxide dismutase. The expression of this recombinant DNA fragment enables to locate the objective protein in mitochondria. Thus the present invention also provides a method for locating an objective protein in mitochondria which comprises expressing the recombinant DNA fragment comprising a coding region for a transit peptide upstream the coding region of an objective protein in an appropriate recombinant host organism.

[0020] As mentioned above, the present invention discloses the nucleotide sequences of active oxygen species-quenching factors, such as mitochondrial superoxide dismutase, cytoplasmic superoxide dismutase and catalase. These polynucleotides are provided for use as probe or primer for cloning gene for the active oxygen species-quenching factor(s) effective in another organism producible of carotenoids on the basis of the homology of the genes.

[0021] FIG. 1 shows the activity staining for superoxide dismutase after native polyacrylamide gel electrophoresis by using cell-free extracts obtained from ATCC 96594 and its *SOD* mutants. Lane 1, *P. rhodozyma* ATCC 96594; Lane 2 *P. rhodozyma* ATCC 96594 :: pSOD/G717 (*SOD1* disruptant); Lane 3 *P. rhodozyma* ATCC 96594 :: pSOD/G828 (*SOD2* disruptant); Lane 4 *P. rhodozyma* ATCC 96594

[0022] As it is briefly explained above, the object of the present invention is to provide a novel process for producing carotenoids biologically. The novel process comprises cultivating a recombinant organism whose gene for one or more active oxygen species-quenching factor(s) is substantially disrupted with the aid of a disruption cassette specific to the said gene, and recovering carotenoids from the culture.

[0023] For the purpose of the present invention, there is also provided a recombinant DNA sequence which contains an open reading frame coding for active oxygen species-quenching factor(s), which may be an enzyme, such as mitochondrial *SOD* or cytoplasmic *SOD*. The recombinant DNA sequence may contain a partial fragment encoding catalase gene. These sequences are useful to construct the said disruption cassette, as they are able to recombine with the native gene(s) for the said enzymes so as to specifically disrupt the gene(s).

[0024] The said recombinant DNA sequence may be the one which can be derived from an organism belonging to the kingdom of *Monera*, *Protista* or *Fungi*, more preferably to the genus *Erwinia*, *Rhodobacter*, *Myxococcus*, *Flavobacter*, *Paracoccus*, *Synechococcus*, *Synechocystis*, *Agrobacterium*, *Streptomyces*, *Haematococcus*, *Dunaliella*, *Phaffia*, *Xanthophyllomyces*, *Neurospora*, *Rhodotorula*, *Blakeslea*, or *Phycomyces*. Particularly preferred organism is *P. rhodozyma*. The active oxygen species-quenching factor coded by the recombinant DNA sequence may be mitochondrial *SOD*, cytoplasmic *SOD* and/or catalase. A specific example of the recombinant DNA sequence is derived from a gene of *Phaffia rhodozyma* and is selected from (i) a DNA sequence represented in SEQ ID NO: 1 or 2; (ii) those cDNAs identified by SEQ ID NO: 4 or 6; (iii) an isocoding or an allelic variant for the DNA sequence represented in SEQ ID NO: 1, 2, 4 or 6; and (iv) a derivative of a DNA sequence represented in SEQ ID NO: 1, 2, 4 or 6 with addition, insertion, deletion and/or substitution of one or more nucleotide(s), and coding for a polypeptide having the said enzyme activity. The said recombinant DNA sequence may also be characterized in that (a) it codes for the said enzyme having an amino acid sequence selected from the group consisting of those described in SEQ ID NOs: 5 and 7, or (b) it codes for a variant of the said enzyme selected from (i) an allelic variant, and (ii) an enzyme having one or more amino acid addition, insertion, deletion and/or substitution and having the stated enzyme activity.

[0025] As used herein, an "allelic variant" means the variant that has at least one mutation in either one of two alleles in the diploid organism such as *Phaffia rhodozyma*, *Xanthophyllomyces dendrorhous* and the like. Both alleles of a given gene are concerned with the same trait or characteristic, but the product or function coded for by a particular allele differs, qualitatively and/or quantitatively, from that coded for by other alleles of that gene.

[0026] Allelic variant can occur naturally or be generated artificially by means of chemical mutagenesis. A wild type allele is one which codes for a particular phenotypic characteristic found in the wild type strain of a given organism.

[0027] As used herein, an "isocoding variant" means the variant in which the nucleotide sequence of a given gene differs from the sequence from the wild type gene although its translated product (*i.e.* amino acid sequence) is identical with that from the wild type protein. This is caused by degeneracy of genetic code and by the difference of codon usage which is not identical among various organisms.

[0028] As used herein, a "derivative of a DNA sequence" is a DNA sequence that encodes a polypeptide having the activity of the corresponding SEQ ID NO but which differs from that DNA sequence by from 1-20, preferably 1-10, such

as 1-5 nucleotide additions, insertions, deletions, and/or substitutions.

[0029] Particularly specified recombinant DNA sequence mentioned above may be that which can be derived from a gene of *Phaffia rhodozyma* and is selected from (i) a DNA sequence represented in SEQ ID NO: 3; (ii) cDNA identified by SEQ ID NO: 8; (iii) an isocoding or an allelic variant for the DNA sequence represented in SEQ ID NO: 3 or 8; and (iv) a derivative of a DNA sequence represented in SEQ ID NO: 3 or 8 with addition, insertion, deletion and/or substitution of one or more nucleotide(s), and coding for a polypeptide having the said enzyme activity. The said recombinant DNA sequence may also be characterized in that (a) it codes for the said enzyme having a partial amino acid sequence selected from the group consisting of those described in SEQ ID NO: 9, or (b) it codes for a variant of the said enzyme selected from (i) an allelic variant, and (ii) an enzyme having one or more amino acid addition, insertion, deletion and/or substitution and having the stated enzyme activity. Such a recombinant DNA sequence may preferably be in the form of a vector.

[0030] The present invention also provides the use of the said recombinant DNA sequence to transform a host organism so as to obtain an organism whose gene for at least one active oxygen species-quenching factor is substantially disrupted with the aid of introduction of the said disruption cassette specific to the said gene.

[0031] As used herein, a gene is "disrupted" or "substantially disrupted" if the activity of the polypeptide that it encodes is reduced relative to a non-disrupted gene. Preferably, the activity is reduced by 10%, preferably by at least 50% such as for example by at least 75%, more preferably by at least 90% to 100%.

[0032] A convenient form of the recombinant DNA sequence may be a vector. The recombinant organism obtained by use of the recombinant DNA is disrupted in its DNA sequence encoding mitochondrial SOD, cytoplasmic SOD, or catalase. The host organism transformed with the recombinant DNA is useful in the improvement of the production process of carotenoids, in particular astaxanthin. Thus the present invention also provides such a recombinant organism.

[0033] This biological production method of carotenoids can improve the productivity of carotenoids, in particular of astaxanthin. Thus the method for producing a carotenoid, characterized in that a recombinant organism described above is cultivated under conditions conducive to the production of the carotenoid is one of the aspect of the present invention. This method may be applied to the biological production of astaxanthin.

[0034] Many researchers pointed out that active oxygen species might stimulate carotenoid production in known carotenogenic organisms. Carotenoid biosynthesis in cyst cells of *Haematococcus pluvialis* is enhanced by environmental oxidative stress (Kobayashi et al., *Appl. Env. Microbiol.*, 59, 867-873, 1993). Carotenoid biosynthesis might be induced by active oxygen species and the accumulated carotenoids might function as a protective agent against oxidative stress damage in *Dunaliella bardawil* (Shaish et al., *Planta*, 190, 363-368, 1993). Although astaxanthin production in *P. rhodozyma* was studied *in vivo* under various cultivation conditions in which a generation of active oxygen species was altered, a correlation between active oxygen generated and carotenoids productivity was not clearly determined, probably because native active oxygen species-quenching factors were still present in such experiments and rescued the effects of active oxygen species on the carotenoids production to some extent (Schroeder, W. A. et al., *J. Gen. Microbiol.*, 139, 907-912, 1993).

[0035] In this invention, to exclude the possibility that existence of native active oxygen species-quenching factor in *P. rhodozyma* can quench the positive effect by active oxygen on astaxanthin production, such native active oxygen species-quenching factors as SOD and catalase were cloned from *P. rhodozyma* to disrupt their expression by constructing and introducing gene disruption plasmids. On the assumption that astaxanthin would play an antioxidant role in *P. rhodozyma*, inactivation of native active oxygen species-quenching factor may affect to carotenoids production probably because relatively increased active oxygen species *in vivo* due to absence of native active oxygen species-quenching factors would stimulate a production of astaxanthin as an alternative player to quench active oxygen species.

[0036] Active oxygen species have toxicity to living cells through the oxidative damage to the intercellular molecules such as proteins or nucleic acids. Recent studies revealed that aging is caused by oxidative damage by demonstrating a correlation between increased superoxide dismutase activity, increased life-span, and decreased oxidative damage in fruit flies and nematodes (Agarwal, S. et al., *Proc. Natl. Acad. Sci. U. S. A.*, 91, 12332-12335, 1994, Larsen, P. L. *Proc. Natl. Acad. Sci. U. S. A.*, 90, 8905-8909, 1993, Sohal, R. S. et al., *J. Biol. Chem.*, 270, 15671-15674, 1995). SOD and related antioxidant enzymes and their genes are studied well in procaryote and eukaryote.

[0037] Yeast, *S. cerevisiae*, like most eucaryotes, contains Cu/ZnSOD (product of the *SOD1* gene) in the cytosol and MnSOD (product of the *SOD2* gene) in the mitochondria. These enzymes catalyze the disproportionating of O_2 , producing O_2 and H_2O_2 . Together with small molecular antioxidants, such as glutathione and ascorbate; other antioxidant enzymes, such as catalases and peroxidases; and metal chelating proteins such as metallothionein, they allow aerobes to survive under O_2 , presumably by minimizing oxidative damage. The importance of cytoplasmic SOD was demonstrated by the high sensitivity to dioxygen shown by *S. cerevisiae* and *Escherichia coli* devoid of SOD. In both organisms, the loss of SOD activity is associated with slow growth in aerobic conditions, with higher mutation rates and with specific biosynthetic defects. (*sod1*⁻ yeast requires lysine and methionine for aerobic growth, whereas *sod*⁻ *E. coli* requires branched amino acids). In some cases these effects are known to be due to the inhibitory effect of

superoxide on iron sulfur cluster proteins (Gardner, P. R. *et al.*, *J. Biol. Chem.*, 266, 19328-19333, 1991, Kuo, C. F., *et al.*, *J. Biol. Chem.*, 262, 4724-4727, 1987). Mutants of *sod2* of *S. cerevisiae* are little affected when grown in air with glucose as the carbon source. However, they are highly sensitive to hyperoxia and grow poorly in normoxia in carbon sources that require respiration for their metabolism.

[0038] Because genes coding for SOD and catalase have been cloned from other species, corresponding genes from *P. rhodozyma* could be cloned with degenerate PCR method. At first, we cloned a partial gene fragment containing a portion of *SOD* gene and *CAT* gene by using above method. The said degenerate PCR is a method to clone a gene of interest which has high homology of amino acid sequence to the known enzyme from other species which has a same or similar function. Degenerate primer, which is used as a primer in degenerate PCR, was designed by a reverse translation of the amino acid sequence to corresponding nucleotides ("degenerated"). In such a degenerate primer, a mixed primer which consists any of A, C, G or T, or a primer containing inosine at an ambiguity code is generally used. In this invention, such the mixed primers were used for degenerate primers to clone above genes. PCR condition used is varied depending on primers and genes to clone as described hereinafter. In this invention, two species of *SOD* genes whose sequence was different from each other were cloned from the same PCR band in degenerate PCR and named as *SOD1* and *SOD2* genes.

[0039] An entire gene containing its coding region with its intron as well as its regulation region such as a promoter or terminator can be cloned from a chromosome by screening of genomic library which is constructed in phage vector or plasmid vector in an appropriate host, by using a partial DNA fragment obtained by degenerate PCR as described above as a probe after it was labeled. Generally, *E. coli* as a host strain and *E. coli* vector, a phage vector such as λ phage vector, or a plasmid vector such as pUC vector are often used in the construction of library and a following genetic manipulation such as a sequencing, a restriction digestion, a ligation and the like. In this invention, an *EcoRI* genomic library of *P. rhodozyma* was constructed in the derivatives of λ vector, λ ZAPII and λ DASHII depending on an insert size. An insert size, what length of insert must be cloned, was determined by the Southern blot hybridization for each gene before a construction of a library. In this invention, a DNA which was used for a probe was labeled with digoxigenin (DIG), a steroid hapten instead of conventional ^{32}P label, following the protocol which was prepared by the supplier (Boehringer-Mannheim (Mannheim, Germany)). A genomic library constructed from the chromosome of *P. rhodozyma* was screened by using a DIG-labeled DNA fragment which had a portion of a gene of interest as a probe. Hybridized plaques were picked up and used for further study. In the case of using λ DASHII (insert size was from 9 kb to 23 kb), prepared λ DNA was digested by the *EcoRI*, followed by the cloning of the *EcoRI* insert into a plasmid vector such as pUC19 or pBluescriptII SK+. When λ ZAPII was used in the construction of the genomic library, *in vivo* excision protocol was conveniently used for the succeeding step of the cloning into the plasmid vector by using a derivative of single stranded M13 phage, Ex assist phage (Stratagene, La Jolla, USA). A plasmid DNA thus obtained was examined for sequencing. In this invention, *SOD1* and *SOD2* genes were obtained from the λ ZAPII library independently from each other and catalase (*CAT*) gene was cloned from λ DASHII library.

[0040] In this invention, we used the automated fluorescent DNA sequencer, ALFred system (Pharmacia, Uppsala, Sweden) using an autocycle sequencing protocol in which the Taq DNA polymerase is employed in most cases of sequencing.

[0041] In this invention, the inventors determined the genomic sequence containing an open reading frame of *SOD1* gene or *SOD2* gene as well as its promoter and terminator sequence. From sequence analysis, it was found that *SOD1* codes for mitochondrial SOD, judged from a presence of transit peptide at its amino terminal end. On the contrary, *SOD2* does not have such a transit peptide sequence and this fact suggested that *SOD2* codes for cytoplasmic SOD. The inventors also determined a partial genomic sequence of an open reading frame for *CAT* gene.

[0042] Transit peptide is a signal sequence to transfer the nucleic gene products, which code on chromosome, and whose translated proteins function in mitochondria, to mitochondrial location of proteins, such as enzymes involved in TCA cycle. To express some proteins in mitochondria, addition of transit peptide at its amino terminal end is useful.

[0043] In this invention, disruption plasmids for *SOD1*, *SOD2* and *CAT* genes were constructed by ligating partial fragments of above genes, which do not contain the either end of genes, with drug resistant genes to suicidal vector, which cannot be autonomously replicated in *P. rhodozyma* due to lack of autonomous replication sequence. A drug resistant gene which encodes the enzyme that enables the host survive in the presence of a toxic antibiotic is often used for the selectable marker. G418 resistance gene harbored in pPR2T (Verdoes *et al.* (International patent publication, WO97/23633)) is an example of a drug resistance gene. Such a suicidal vector cannot replicate by itself and can be present only in an integrated form on the chromosome of the host as a result of a single-crossing recombination using the homologous sequence between a vector and a chromosome. In the case of recombination with a gene of interest, its genetic sequence cannot be reconstituted on the chromosome of the host strain due to lack of either end of the gene and as a consequence, the gene of interest could be disrupted in the recombinant strain thus obtained. Another example for disruption plasmid is a double crossing over type of plasmid. This type of disruption plasmid contains two different partial fragments of objective gene to be disrupted and selective marker gene such as a drug resistant gene is inserted between such two fragments. After the recombination between the chromosome of the re-

recipient cells and donor plasmid DNA at the two homologous parts of the gene, replacement of the chromosome sequence with the donor DNA occur and selective marker gene is inserted into the objective gene to be disrupted. In general, double crossing over type of plasmid has lower frequency of recombination than single crossing type of vector.

[0044] In this invention, inventors inactivated the enzymes of interest by their disruption of corresponding genes. The other way to evaluate an effect of gene product of an interest is to decrease its expression with genetic engineering methods. For this purpose, some methods were exploited. One of such methods is anti-sense method. Anti-sense method is a method to decrease an expression of gene of interest by introducing an artificial gene fragment, whose sequence is complementary to that of gene of interest. And such an anti-sense gene fragment form complex with mature mRNA fragment of objective gene *in vivo* and inhibit an efficient translation from mRNA, as a consequence.

[0045] The other method is a mutation of promoter region. In general, the gene consists of several parts which have different functions from each other. In eukaryotes, genes which encode corresponding protein are transcribed to pre-mature messenger RNA (pre-mRNA) differing from the genes for ribosomal RNA (rRNA), small nuclear RNA (snRNA) and transfer RNA (tRNA). Although RNA polymerase II (PolII) plays a central role in this transcription event, PolII can not solely start transcription without *cis* element covering an upstream region containing a promoter and an upstream activation sequence (UAS), and a *trans*-acting protein factor. At first, a transcription initiation complex which consists of several basic protein components recognize the promoter sequence in the 5'-adjacent region of the gene to be expressed. In this event, some additional participants are required in the case of the gene which is expressed under some specific regulation, such as a heat shock response, or adaptation to a nutrition starvation, and so on. In such a case, a UAS is required to exist in the 5'-untranslated upstream region around the promoter sequence, and some positive or negative regulator proteins recognize and bind to the UAS. The strength of the binding of transcription initiation complex to the promoter sequence is affected by such a binding of the *trans*-acting factor around the promoter, and this enables the regulation of the transcription activity.

[0046] After the activation of a transcription initiation complex by the phosphorylation, a transcription initiation complex initiates transcription from the transcription start site. Some parts of the transcription initiation complex are detached as an elongation complex from the promoter region to the 3' direction of the gene (this step is called as a promoter clearance event) and an elongation complex continues the transcription until it reaches to a termination sequence that is located in the 3'-adjacent downstream region of the gene.

[0047] To decrease an expression of gene of interest, mutation by conventional chemical mutagenesis or genetic site-directed mutagenesis in the promoter region of objective gene containing UAS sequence described above was often used. Mutant strain in which expression of enzyme of interest might decrease can be obtained by transforming host strain with a recombinant DNA having such a mutated promoter region. As described above, such an attempt to decrease the expression of gene is also employed as well as gene disruption for objective to determine an effect of gene product on phenomena in living organisms.

[0048] As a transformation method, LiCl method (Wery *et al.*, *Yeast*, 12 (7), 641-651, 1996) and electroporation method (Wery *et al.*, *Gene*, 184, 89-97, 1997) were applied to transform *P. rhodozyma* but the efficiency of transformation under these conditions seemed to be still low. In this invention, Biolistic transformation method (Johnston *et al.*, *Methods in Molecular Biology*, 53; 147-153, 1996) was applied for transformation of *P. rhodozyma*. Biolistic method is a simple and reliable protocol, in which donor DNA coated on gold or tungsten particle is shot into the living cells directly with high-pressured helium gas. This transformation protocol was successfully applied to *Cryptococcus neoformans* which belongs to basidiomycetous yeast as well as *P. rhodozyma* and was difficult to transform with conventional transformation method (Toffaletti, *et al.*, *J. Bacteriol.*, 175 (5), 1405-1411, 1993). In this invention, this biolistic method was successfully used to transform *P. rhodozyma* cells.

[0049] The event of gene disruption can be confirmed by enzymatic characterization directly and by genetic analysis with PCR or Southern blot hybridization by using the chromosome obtained from transformants thus obtained. In this invention, a direct confirmation of SOD disruption were performed by activity staining and characterization of catalase disruption were conducted by visual observation like catalase test which is often used in bacterial taxonomy.

[0050] Such a genetically engineered *P. rhodozyma* would be cultivated in an appropriate medium and evaluated in its productivity of astaxanthin.

[0051] The present invention is further illustrated with Examples described below by referring to the attached drawings.

Example

[0052] The following materials and methods were employed in the Example described below:

Strain**[0053]**

- 5 *P. rhodozyma* ATCC 96594 (re-deposited under the accession No. ATCC 74438 on April 8, 1998 pursuant to the Budapest Treaty)
- E. coli* DH5a: F⁻, ϕ 80d, *lacZ* Δ M15, Δ (*lacZYA-argF*)U169, *hsd* (*r*_K⁻, *m*_K⁺), *recA1*, *endA1*, *deoR*, *thi-1*, *supE44*, *gyrA96*, *relA1* (Toyobo, Osaka, Japan)
- 10 *E. coli* XL 1 -Blue MRF': Δ (*mcrA*)183, Δ (*mcrCB-hsdSMR-mrr*)173, *endA1*, *supE44*, *thi-1*, *recA1*, *gyrA96*, *relA1*, *lac* [F' *proAB*, *lac1* Δ M15, Tn10 (*tet*^r)] (Stratagene, La Jolla, USA)
- 15 *E. coli* SOLR: *e14'* (*mcrA*), Δ (*mcrCB-hsdSMR-mrr*)171, *sbcC*, *recB*, *recJ*, *umuC* :: Tn5(*kan*^r), *uvrC*, *lac*, *gyrA96*, *relA1*, *thi-1*, *endA1*, λ ^R, [F' *proAB*, *lac* Δ M15] Su⁻ (nonsuppressing) (Stratagene)
- E. coli* XL1/MRA (P2): Δ (*mcrA*)183, Δ (*mcrCB-hsdSMR-mrr*)173, *endA1*, *supE44*, *thi-1*, *gyrA96*, *relA1*, *lac* (P2 lysogen) (Stratagene)
- 20 *E. coli* TOP10: F⁻, *mcrA*, Δ (*mrr-hsdRMS-mcr*)BC, ϕ 80, Δ *lacZ* M15, Δ *lacX*74, *recA1*, *deoR*, *araD139*, (*ara-leu*)7697, *galU*, *galK*, *rpsL* (*Str*^r), *endA1*, *nupG* (Invitrogen, Carlsbad, USA)

Vector**[0054]**

- λ ZAPII (Stratagene)
- λ DASHII (Stratagene)
- 30 pBluescriptII SK+ (Stratagene)
- pCR2.1TOPO (Invitrogen)
- 35 pUC19 (Takara Shuzo, Otsu, Japan)

Media

- 40 **[0055]** *P. rhodozyma* strain is maintained routinely on agar plate of YPD medium (DIFCO, Detroit, USA). *E. coli* strain is maintained in LB medium (10 g Bacto-trypton, 5 g yeast extract (DIFCO) and 5 g NaCl per liter). NZY medium (5 g NaCl, 2 g MgSO₄·7H₂O, 5 g yeast extract (DIFCO), 10 g NZ amine type A (WAKO, Osaka, Japan) per liter is used for λ phage propagation in a soft agar (0.7 % agar (WAKO)). When an agar medium was prepared, 1.5 % of agar (WAKO) was supplemented.

[0056] Methods

- [0056]** A general method for techniques of molecular genetics was according to the method in Molecular cloning: a Laboratory Manual, 2nd Edition (Cold Spring Harbor Laboratory Press, 1989). Restriction enzymes and T4 DNA ligase were purchased from Takara Shuzo (Japan).
- 50 **[0057]** Isolation of a chromosomal DNA from *P. rhodozyma* was performed by using QIAGEN Genomic Kit (QIAGEN, Hilden, Germany) following the protocol supplied by the manufacturer. Mini-prep of plasmid DNA from transformed *E. coli* was performed with the Automatic DNA isolation system (PI-50, Kurabo, Co. Ltd., Osaka, Japan). Midi-prep of plasmid DNA from an *E. coli* transformant was performed by using QIAGEN column (QIAGEN). Isolation of λ DNA was performed by Wizard lambda preps DNA purification system (Promega, Madison, USA) following the protocol prepared by the manufacturer. A DNA fragment was isolated and purified from agarose by using QIAquick or QIAEX II (QIAGEN).
- 55 **[0058]** Isolation of total RNA from *P. rhodozyma* was performed with phenol method by using Isogen (Nippon Gene, Toyama, Japan). mRNA was purified from total RNA thus obtained by using mRNA separation kit (Clontech, Palo Alto,

USA). cDNA was synthesized by using CapFinder cDNA construction kit (Clontech).

[0059] *In vitro* packaging was performed by using Gigapack III gold packaging extract (Stratagene).

[0060] Polymerase chain reaction (PCR) is performed with the thermal cycler from Perkin Elmer model 2400. Each PCR condition is described in examples. PCR primers were purchased from a commercial supplier or synthesized with a DNA synthesizer (model 392, Perkin Elmer, Japan, Urayasu, Japan). Fluorescent DNA primers for DNA sequencing were purchased from Pharmacia. DNA sequencing was performed with the automated fluorescent DNA sequencer (ALFred, Pharmacia).

[0061] Competent cells of *E. coli* DH5 α were purchased from Toyobo (Japan).

[0062] The apparatus and reagent for biolistic transformation of *P. rhodozyma* were purchased from Nippon Bio-Rad Laboratories (Tokyo, Japan).

Example 1: Isolation of mRNA from *P. rhodozyma* and construction of cDNA library

[0063] To construct cDNA library of *P. rhodozyma*, total RNA was isolated by phenol extraction method right after the cell disruption and the mRNA from *P. rhodozyma* ATCC96594 strain was purified by using mRNA separation kit (Clontech).

[0064] At first, Cells of ATCC96594 strain from 10 ml of two-day-culture in YPD medium were harvested by centrifugation (1500 x g for 10 min.) and washed once with extraction buffer (10 mM Na-citrate / HCl (pH 6.2) containing 0.7 M KCl). After suspending in 2.5 ml of extraction buffer, the cells were disrupted by French press homogenizer (Ohtake Works Corp., Tokyo, Japan) at 1500 kgf/cm² and immediately mixed with two times of volume of isogen (Nippon gene) according to the method specified by the manufacturer. In this step, 400 μ g of total RNA was recovered.

[0065] Then this total RNA was purified by using mRNA separation kit (Clontech) according to the method specified by the manufacturer. Finally, 16 μ g of mRNA from *P. rhodozyma* ATCC 96594 strain was obtained.

[0066] To obtain cDNA species from *P. rhodozyma*, CapFinder PCR cDNA construction kit (Clontech) was used according to the method specified by the manufacturer. One μ g of purified mRNA was applied for a first strand synthesis followed by PCR amplification. After this amplification by PCR, 1 mg of cDNA pool was obtained.

Example 2: Cloning of two species of partial SOD gene from *P. rhodozyma*

[0067] To clone a partial SOD gene from *P. rhodozyma*, a degenerate PCR method was exploited. Two mixed primers whose nucleotide sequences were designed and synthesized as shown in TABLE 1 based on the common sequence of known superoxide dismutase genes from other species.

TABLE 1

Sequence of primers used in the cloning of SOD1 and SOD2 genes

Sod1 ; AARCAYCAYCARACNTAYGTNAA (sense primer) (SEQ ID NO: 10)

Sod4 ; GCCCANCCNGANCCYTGNACNCC (antisense primer) (SEQ ID NO: 11)

(R=A or G; Y=C or T; N=A, C, G or T)

[0068] After the PCR reaction of 25 cycles of 94 °C for 15 seconds; 46 °C for 30 seconds and 72°C for 15 seconds by using ExTaq (Takara Shuzo) as a DNA polymerase and cDNA pool obtained in example 1 as a template, reaction mixture was applied to agarose gel electrophoresis. A PCR band that has a desired length was recovered and purified by QIAquick (QIAGEN) according to the method by the manufacturer and then ligated to pCR2.1-TOPO (Invitrogen). After the transformation of competent *E. coli* TOP10, 6 white colonies were selected and plasmids were isolated with Automatic DNA isolation system (Kurabo PI-50). As a result of sequencing, it was found that two clones had different sequences each other, whose deduced amino acid sequences were independently similar to known SOD genes. These isolated cDNA clones were designated as pSOD614 #2 and pSOD614 #3 and used for further study.

Example 3: Isolation of genomic DNA from *P. rhodozyma*

[0069] To isolate a genomic DNA from *P. rhodozyma*, QIAGEN genomic kit was used according to the method spec-

ified by the manufacturer.

[0070] At first, cells of *P. rhodozyma* ATCC 96594 strain from 100 ml of overnight culture in YPD medium were harvested by centrifugation (1500 x g for 10 min.) and washed once with TE buffer (10 mM Tris / HCl (pH 8.0) containing 1 mM EDTA). After suspending in 8 ml of Y1 buffer of the QIAGEN genomic kit, lyticase (SIGMA) was added at the concentration of 2 mg/ml to disrupt cells by enzymatic degradation and the reaction mixture was incubated for 90 minutes at 30 °C and then proceeded to the next extraction step. Finally, 20 µg of genomic DNA was obtained.

Example 4: Southern blot hybridization by using pSOD614 #2 and pSOD614 #3 as probes

[0071] Southern blot hybridization was performed to clone genomic fragment which contains *SOD* genes from *P. rhodozyma*. Two µg of genomic DNA was digested by *EcoRI* and subjected to agarose gel electrophoresis followed by acidic and alkaline treatment. The denatured DNA was transferred to nylon membrane (Hybond N+, Amersham) by using transblot (Joto Rika, Tokyo, Japan) for an hour. The DNA was transferred to nylon membrane and fixed by a heat treatment (80 °C, 90 minutes). Probes were prepared by labeling the template DNAs (*EcoRI*- digested pSOD614 #2 and pSOD614 #3) with DIG multipriming method (Boehringer Mannheim). Hybridization was performed with the method specified by the manufacturer. As a result, hybridized bands were visualized in the length of 7.5 kilobases (kb) against the probe prepared from pSOD614 #2, and in the length of 8.0 kilobases (kb) against the probe prepared from pSOD614 #3.

Example 5: Cloning of genomic fragments containing *SOD* genes whose sequences were different from each other

[0072] Four µg of the genomic DNA was digested by *EcoRI* and subjected to agarose gel electrophoresis. Then, DNAs whose length are within the range from 7 to 9 kb were recovered by a conventional elution method by using dialysis membrane. The purified DNA was ligated to 1 µg of *EcoRI*-digested and CIAP (calf intestine alkaline phosphatase) - treated λZAPII (Stratagene) at 16 °C overnight, and packaged with Gigapack III gold packaging extract (Stratagene). The packaged extract was infected to *E. coli* XL1Blue MRF' strain and over-laid with NZY medium poured onto LB agar medium. About 6000 plaques were screened by using *EcoRI*- digested pSOD614 #2 and pSOD614 #3 as probes. Six plaques were hybridized to the labeled pSOD614 #2 probe and two plaques were hybridized to the labeled pSOD614 #3 probe. Then, hybridized plaques were subjected to *in vivo* excision protocol according to the method specified by the manufacturer (Stratagene). It was found that one plasmid isolated from six of pSOD614 #2-hybridized plaques had the same fragments as that of pSOD614 #2 as a result of PCR analysis by using *sod1* and *sod4* primers. This plasmid was named as pSOD703. It was also found that two plasmids isolated from two of pSOD614 #3-hybridized plaques had the same fragment as that of pSOD614 #3 as a result of PCR analysis by using *sod1* and *sod4* primers. One of the plasmids was named as pSOD626 and used for further study.

Example 6: Sequence analysis of two species of MnSOD genes obtained from *P. rhodozyma*

[0073] A complete nucleotide sequence was determined by sequencing of pSOD703 and pSOD626 with a primer-walking procedure. Nucleotide and their deduced amino acid sequences for *SOD1* gene which harbored on pSOD703 and for *SOD2* gene which harbored on pSOD626 are listed as SEQ ID NO: 1, NO: 2, NO: 5 and NO: 7 in the sequence listing section.

[0074] Both of deduced amino acid sequences of *SOD1* and *SOD2* genes were homologous to known MnSODs obtained from other species and not to Cu/ZnSODs or FeSOD as a result of BLAST analysis (Altschul, S.F. *et al.*, *J. Mol. Biol.* 215, 403-410, 1990).

[0075] *SOD1* gene had 7 introns and 8 exons. And its deduced open reading frame consisted of 223 amino acids. On the other hand, *SOD2* gene had 10 introns and 11 exons, and its deduced open reading frame consisted of 199 amino acids. Most difference between two isolated *SOD* genes was extended region of *SOD1* gene in its amino terminal end, whose sequence might act as a transit peptide to mitochondria

[0076] In fact, Schroeder *et al.* reported two species of SODs which were detected as KCN and H₂O₂ resistant SODs in activity staining of native polyacrylamide gel electrophoresis (PAGE) in *P. rhodozyma*. They commented two species of MnSOD were indicated as aggregates or isozymes and did not refer to their precise nature and their subcellular location. As described in the following section, it was clarified that two species of those KCN- and H₂O₂- resistant SODs (i.e. MnSOD) were products of *SOD1* and *SOD2* genes. As described in the section of "detailed description of the invention", most eucaryotes have different species of SODs in their intracellular location (MnSOD in the mitochondrial fraction and Cu/ZnSOD in the cytoplasmic fraction). This is the first example in which two species of MnSODs functioned in a different subcellular location.

Example 7: Construction of disruption plasmids for *SOD1* and *SOD2* genes

[0077] As described in the section "detailed description of the invention", a plasmid harboring a drug resistant marker cassette was constructed by inserting G418 resistant structure gene between promoter and terminator genes of glyceraldehyde-3-phosphate dehydrogenase (GAP) and ligating into *KpnI*- and *HindIII*- digested pUC19. This plasmid was named as pUC-G418 and used for a further study. As gene fragments which were used for homologous recombination, partial fragments of *SOD1* and *SOD2* genes were synthesized *in vitro* by PCR method using PCR primers whose sequences were shown in TABLE 2.

TABLE 2

Sod14 ; GGTACCTCCGATGATAGGAATGTGAG (sense primer) (SEQ ID NO: 12)

Sod15 ; GAATTCAGTTCAACGGAGGAGGACAC (antisense primer) (SEQ ID NO: 13)

Sod47; GAATTCGGAGGAGGACACATCAACCG (sense primer) (SEQ ID NO: 14)

Sod48; GGTACCTGTACTGGAGGTAGAAAGCG (antisense primer) (SEQ ID NO: 15)

[0078] . PCR condition was as follows; 25 cycles of 94 °C for 15 seconds, 50 °C for 30 seconds and 72 °C for 15 minutes. As a template, 0.1 µg of genomic DNA obtained in Example 3 was used, and ExTaq as a DNA polymerase. A partial fragment of *SOD1* which could be obtained from PCR using sod14 and sod15 as primers and a partial *SOD2* gene which could be obtained from PCR using sod47 and sod48 as primers were amplified respectively. Each amplified 0.65 kb fragments was recovered and cloned in pCR2.1-TOPO (Invitrogen) according to the protocol specified by the manufacturer. Six independent clones from white colonies of *E. coli* TOP10 transformants were selected and plasmids were prepared from those transformants. As a result of restriction analysis and sequencing, one clone which had a partial *SOD1* gene was selected for a further study and named as pSOD715. In a similar manner, one clone which had a partial *SOD2* gene was selected and named as pSOD826.

[0079] Then, pSOD715 and pSOD826 were digested by *EcoRI* and *KpnI*, and 0.65kb fragments yielded were purified by using QIAquick protocol and ligated to *EcoRI*- and *KpnI*- digested pUC19-G418. Six independent clones from ampicillin-resistant colonies of *E. coli* DH5α transformants were selected and plasmids were prepared from those transformants. As a result of restriction analysis and sequencing, one clone in which a partial *SOD1* was fused to G418 resistant cassette was obtained and named as pSOD/G717. In a similar manner, one clone in which a partial *SOD2* was fused to G418 resistant cassette was obtained and named as pSOD/G828.

Example 8: Transformation of *P. rhodozyma* ATCC96594 with biolistic method

[0080] Transformation protocol were followed to the method described in Methods in Molecular Biology (Johnston *et al.*, 53; 147-153, 1996). As a host strain, *P. rhodozyma* ATCC 96594 was cultured in YPD medium to stationary phase. After centrifugation of broth, cells were concentrated by 10-fold with sterilized water and 200 µl of cell suspension were spread on YPD medium containing 100 µg of geneticin, and 0.75 M of mannitol and sorbitol. Five micrograms of circular DNA of pSOD/G717 and pSOD/G828 were coated on 1.5 mg of 0.9 µm gold particle, and used as donor DNA for biolistic transformation. Hundreds of geneticin-resistant clones were yielded after one week of incubation at 20 °C. Four of those transformants were selected and chromosomes were prepared from them. One of the transformants was confirmed to have disrupted structure of chromosomal *SOD1* or *SOD2* gene by PCR and Southern blot hybridization analyses, and was used for further study.

Example 9: Activity staining of native PAGE by using crude extracts which obtained from candidates for *SOD1* and *SOD2* disruptants

[0081] ATCC96594 strain and candidates which obtained from biolistic transformation of ATCC96594 were cultivated in YPD medium for two days and harvested by centrifugation for 10 min. by 3000 x g at 4 °C. After wash with Tris-HCl buffer (10mM / pH 8.0), cells were concentrated by 10-fold with the same buffer. Cells were disrupted with French

press homogenizer (Ohtake Works) at 1500 kg/cm² and crude extract was prepared after a microcentrifugation at 15000 rpm (TOMY, MRX150) of homogenized fraction.

[0082] Protein concentration of crude extract thus prepared was determined with BCA protein assay reagent released by PIERCE (Rockford, U.S.A.). A volume of crude extract corresponding to 300 µg of protein were subjected to native PAGE according to the method described by Schroeder W. A. *et al.* (*J. Gen. Microbiol.*, 139, 907-912, 1993). Method of activity staining was referred to the method by Flohé *et al.* (*Methods in Enzymology*, 105, 93-104, 1984).

[0083] Result of activity staining is depicted in FIG. 1. In the extract of parental strain, ATCC96594, two bands were visualized as transparent bands which have SOD activity in the dark background. On the contrary, ATCC96594 :: pSOD/G717 strain in which *SOD1* gene was disrupted, lacked activity band with high mobility in native PAGE and ATCC96594 :: pSOD/G828 strain, in which *SOD2* gene was disrupted lacks activity band with low mobility in native PAGE. From this result, it was found that two species of MnSOD that were present in the crude extract of *P. rhodozyma* were products of *SOD1* and *SOD2* genes, and SOD species with high mobility and with low mobility in native PAGE corresponded to *SOD1* and *SOD2* gene product, respectively.

Example 10: Cloning of partial catalase (*CAT*) gene from *P. rhodozyma*

[0084] To clone a partial *CAT* gene from *P. rhodozyma*, a degenerate PCR method was exploited. Two mixed primers whose nucleotide sequences were designed and synthesized as shown in TABLE 3 based on the common sequence of known catalase genes from other species.

TABLE 3

Sequence of primers used in the cloning of *CAT* genes

Cat2 ; MIGNTTYTCNACNGTNGGNGGNGA (sense primer) (SEQ ID NO: 16)

Cat5 ; CKRTGNCKYTGNGTRTCNGGRTA (antisense primer) (SEQ ID NO: 17)
(M=A or C; N=A, C, G or T; Y=C or T; K=G or T; R=A or G)

[0085] After the PCR reaction of 25 cycles of 94 °C for 15 seconds, 45 °C for 30 seconds and 72°C for 15 seconds by using ExTaq (Takara Shuzo) as a DNA polymerase and genomic DNA obtained in example 3 as a template, reaction mixture was applied to agarose gel electrophoresis. A PCR band that has a 1.0 kb length was recovered and purified by QIAquick (QIAGEN) according to the method by the manufacturer and then ligated to pCR2.1-TOPO (Invitrogen). After the transformation of competent *E. coli* TOP 10, 6 white colonies were selected and plasmids were isolated with Automatic DNA isolation system. As a result of sequencing, it was found that two clones had sequence whose deduced amino acid sequences was similar to known *CAT* genes. One of these isolated DNA clones was designated as pCAT702 and used for further study.

Example 1 Cloning of genomic fragments containing *CAT* gene

[0086] In a similar manner to Example 4, Southern blot hybridization study was performed by using pCAT702 as a probe. As a result, hybridized band which had a size from 9 kb to 23 kb was visualized. Next, 4 µg of the genomic DNA was digested by *EcoRI* and subjected to agarose gel electrophoresis. Then, DNAs whose length is within the range from 9 to 23 kb was recovered by a conventional elution method by using dialysis membrane. The purified DNA was ligated to 1 µg of *EcoRI*-digested and CIAP (calf intestine alkaline phosphatase) - treated λDASHII (Stratagene) at 16 °C overnight, and packaged by Gigapack III gold packaging extract (Stratagene). The packaged extract was infected to *E. coli* XL1Blue MRA(P2) strain and over-laid with NZY medium poured onto LB agar medium. About 8000 plaques were screened using *EcoRI*- digested pCAT702 as a probe. Six plaques were hybridized to the labeled pCAT702 probe. λDNA was prepared from each λ clone and it was found that 4 of 6 clones contained same fragment to the insert of pCAT702 as a result of PCR using Cat2 and Cat5 primers, and sequencing analysis. A partial nucleotide and its deduced amino acid sequence for *CAT* gene are listed as SEQID NO: 3 and SEQID NO: 9 in the sequence listing section.

Example 12 Construction of disruption plasmid for *CAT* gene

[0087] In a similar manner to Example 7, disruption plasmid for *CAT* gene was constructed. At first, *SacI* linker was

inserted at *Hind*III site of pUC19-G418, in which terminator region of G418-resistant cassette was located, and as a result of restriction analysis, pUC19-G418Sa, which had *Sac*I site at the end of the G418-resistant cassette was yielded. Then, *Kpn*I- and *Sac*I- fragment derived from pUC19-G419Sa was ligated to *Kpn*I- and *Sac*I- digested pCAT702 and yielded pCAT/G706, in which a partial genomic *CAT* gene was fused to G418-resistant cassette.

Example 13 Transformation of *P. rhodozyma* ATCC 96594 by using pCAT/G706 as donor plasmid

[0088] In a same manner to Example 8, *P. rhodozyma* ATCC 96594 was transformed with *CAT*-disruption plasmid, pCAT/G706. Hundreds of geneticin-resistant clones were yielded after one week of incubation at 20 °C. Four of those transformants were selected and chromosomes were prepared from them. One of the transformants was confirmed to have disrupted structure of chromosomal *CAT* gene by PCR and Southern blot hybridization analyses, and was used for further study.

[0089] Then, two candidates for *CAT* disruptant were characterized with catalase test, which was often used in bacterial taxonomic study. One loopful of cells of *P. rhodozyma* was soaked in 3% H₂O₂ solution and then occurrence of dioxygen gas was observed. Although an immediate occurrence of O₂ foam were confirmed when ATCC 96594 cells were applied to this catalase test, O₂ foam was occurred after a long lag when two ATCC 96594 :: pCAT/G706 mutants were soaked in H₂O₂ solution. From this result, disruption of *CAT* gene was suggested, but a remaining weak activity indicated a presence of another player who catalyzes disappearance of H₂O₂ such as peroxidase in *P. rhodozyma*.

Example 14 Evaluation of *SOD1*, *SOD2* and *CAT* disruptants derived from *P. rhodozyma* for their astaxanthin production

[0090] Effect of gene disruption of *SOD1*, *SOD2* and *CAT* gene on astaxanthin production was evaluated by cultivation in YPD medium with shaking flasks. Cells which grew on YPD agar were suspended in YPD medium and a portion of cell suspension was inoculated to 50 ml of YPD medium in 500 ml baffled flask. Cells were grown with 200 r. p. m. at 20 °C for 84 hours. At an appropriate interval, 3 ml of broth was withdrawn and was analyzed for cell yield, consumption of glucose and astaxanthin content.

[0091] Cell yield was measured as optical density at 660 nm and as dry cell weight by weighing cells after filtration through 0.45 µm cellulose acetate plus nitrocellulose membrane (HAWP04700, Millipore, Bedford, U.S.A.) and heating at 80 °C overnight. Astaxanthin content of *P. rhodozyma* was measured with HPLC method after extraction of carotenoids from cells of *P. rhodozyma* by disruption with glass beads. After extraction, 5ml of acetone/BHT/water containing appropriate concentration of bixin as an internal standard was added. Supernatant was analyzed for astaxanthin content with following HPLC system.

HPLC column; YMC-Pak ODS-A (6 mm, 150 mm)

Temperature; room temperature

Eluent; acetonitrile / methanol / isopropanol (85/10/5)

Injection volume; 10 µl

Flow Rate; 2.0 ml/minute

Detection; UV at 471 nm

Results obtained from 84 hour-culture are summarized in TABLE 4.

TABLE 4

The effect of <i>SOD</i> and <i>CAT</i> mutation on productivity of total carotenoids and astaxanthin by <i>P. rhodozyma</i>		
Strain	Total carotenoids (mg/g-dry-cell)	astaxanthin (mg/g-dry cell)
ATCC 96594	0.169	0.111
Δ <i>SOD1</i>	0.259	0.146
Δ <i>SOD2</i>	0.202	0.129
Δ <i>CAT</i>	0.229	0.144

[0092] *SOD1* and *SOD2* disruptants showed elevated level of productivity for total carotenoids as well as astaxanthin compared to their host strain, ATCC 96594. Especially, *SOD1* disruptant showed significant increase of carotenoids and astaxanthin production by 53.3 % and 31.5 %, respectively.

[0093] *SOD1* seemed to be a mitochondrial enzyme judged from deduced transit peptide sequence at its amino terminal end and might act to scavenge superoxide radical, a kind of active oxygen species occurred in the respiratory chain at mitochondria. These data suggested that astaxanthin production was stimulated by a generation of intracellular active oxygen to compensate the lack of native player of active oxygen species-quenching factor, *SOD1*.

5

10

15

20

25

30

35

40

45

50

55

SEQUENCE LISTING

<110> F. HOFFMANN-LA ROCHE AG

5 <120> Improvement of biological production of carotenoids and
biological materials therefor

<130> SOD

10 <140>
<141>

<150> EP99123821.3
<151> 1999-12-01

15 <160> 17

<170> PatentIn Ver. 2.0

20 <210> 1
<211> 3632
<212> DNA
<213> *Phaffia rhodozyma*

<220>
<221> 5'UTR
<222> (922)..(923)
<223> EXPERIMENTAL

25 <220>
<221> exon
<222> (986)..(1096)

30 <220>
<221> intron
<222> (1097)..(1306)

<220>
<221> exon
<222> (1307)..(1456)

35 <220>
<221> intron
<222> (1457)..(1555)

40 <220>
<221> exon
<222> (1556)..(1589)

<220>
<221> intron
<222> (1590)..(1694)

45 <220>
<221> exon
<222> (1695)..(1799)

<220>
<221> intron
<222> (1800)..(1920)

50 <220>
<221> exon
<222> (1921)..(1982)

55 <220>
<221> intron
<222> (1983)..(2076)

EP 1 111 067 A2

```

<220>
<221> exon
<222> (2077)..(2140)

5
<220>
<221> intron
<222> (2141)..(2246)

<220>
<221> exon
10 <222> (2247)..(2272)

<220>
<221> intron
<222> (2273)..(2390)

15
<220>
<221> exon
<222> (2391)..(2507)

<220>
<221> polyA.site
20 <222> (2663)..(2664)
<223> EXPERIMENTAL

<400> 1
tccgtgtgat aatcttttcta acgccttgta ctttgaccaa ggcgtttgtc cgaaattttg 60
25 caaacttagt gtttggtcga tggacgggtct tcggtaccag aactgacggc tcgccaataa 120
agtatgacga tggtagaggt gaaggaggga accacagggt gaccagtctc aaagagtgtc 180
gatgtgcgcg aggtatttgc attaaatggg gttgtatatg ctagagccaa gagaagacat 240
30 ttggttttgg ttttggtttt gcatttgatg agatgtgtca cgattgaaga cgggaggagg 300
ctcactaacc caagaagcca ggatcaggag gaatgcctcc cccttttcat caagatcttt 360
ctcacatcga acatttgaca ttctcttttag tatecttcta tccttttctt ccaacttctc 420
35 ccattgtatc gaatttgctc gaattgctct tcttatctct gagcagagat gggcattcca 480
atatcgaagg agcgacacaa gaccttgagg tttgggtaac agatgaagag gggccgagggt 540
ggatgggggt gtaggaagta gctgatcgat gagttcctgg atgatgatag gcgaagggaac 600
40 agacatagga tctctgtctc gtcttggaat tactgagtct tgtatccagc gtgttcttgt 660
ctcgaagaag ccttcaagat cgatgtaaga taagacaggc aatgaggacg gacgaacgaa 720
cgaacgaaaa gaacagaaga gctggtaagt cagtcagtca gtcagtcagt caatcaaaca 780
45 ctggtgtcta gggttatagc tcgacgcgac gcgacgcgtt tgagacgca tatgcttacg 840
taatacctgg cgicacccc ccagccgagg caagagccga gccgctctgt aacgacaaaa 900
ttcaaaaggc tttctccatc ttaagctcat tctcatctaa ccgactcacc tcgttcccat 960
50 cattccacc atctaccgc catcc atg tct gtt cga gca tcc ctg tct tcc 1012
gtg tct aga cag act ttc gtc gct cct gct gct ttc cag atc agg gca 1060
aag cat acc ctg cct gag ctt cct tac gct tac gat gtaagacttt 1106
55 tccgtgttct cctattcgtc gctttcttgg ttttttctgt cttcgccctc tagctcttct 1166

```

EP 1 111 067 A2

tcgtcccttc tgctctgctc tttgttggtg atattcagct cgatagacta acccatctca 1226
 tctctggac attcttttac tggaaacgta tcttgctctt gggttttctt ggctttgggt 1286
 5 gaaaattcct ctccactcag gcc ctg gag ccc tcc atc tcc aag gag atc atg 1339
 acc ctt cac cac acc aag cac cat cag act tat gtt aac ggc ctg aac 1387
 gct gcc gag gag agc tac tgg gcc gct gtg ggc aag gag gat gtg ctt 1435
 10 acc cag gtt aag ctt cag tct gtacgtctga cctgtttttt atcgaccgga 1486
 acgcttggtg agggaggaga tgaagtttga tgagcgctca tcgtctagca cgttgaccgg 1546
 atcatacag gct ctg aag ttc aac gga gga gga cac atc aat c 1589
 15 gtcagtgata tcttcaaac tcttgctgag caagtcaggt caagctgact gtttcgcttt 1649
 gttctgagg atctatctca tcttgattt ggcatgatga aacag-ac tct ctg ttc 1705
 tgg aag aac ttg gct ccc tat gga tcc gag gag gct acc ctg tct gaa 1753
 20 gga cct ctg aag aag gct atc gag gaa tct ttt ggt tct ttc gag g 1799
 gtccgtccat ctatcttctt attcagttgt gtttggttcc ggtatactca tctgttttgt 1859
 tccccacaa aataaaaaata aaaatcttgt cctctccggg ggctcgactg caggttcata 1919
 25 g cc ttc aag aag aag ttc aac gct gac acc gct gct gtc caa gga tcc 1967
 gga tgg ggc tgg ctt gtatgtatca tctctttcc atctcaaact ctctcagag 2022
 tctttttctt tgagacttca aactgactat acatgtttct acaacaaaca acag ggc 2079
 30 ttg aac ccg ctt act aag aag ctg gaa gtc acc acg acc gcc aac cag 2127
 gac cct ctg ctt a gtaagttgt tctacatgat tttctatctc aacggatct 2180
 gcatgatctg tcaatgatcc actggattct cttgtttcgt tttctcggg atgatttcac 2240
 35 aaacag ct cac att cct atc atc gga gtt gac gtgcgtatct tcttgaata 2292
 gtcgtagcgt ctgatctcgt tttattgact gacgtgttgc ttctgtccaa atcattaaaa 2352
 aaaatgaaaa caaataatcg attgaccgac gaaacag atc tgg gag cac gct ttc 2408
 40 tac ctt cag tac aag aac gtc aag cct gac tat ctg gct gct gtt tgg 2456
 tcc gtt atc aac tac aag gag gca gag gcc cga ttg cag gct gct ctg 2504
 taa gcgggacgaa aagtaacgac atatgaaggg aggatcaaat atcgtttctt 2557
 45 cataaacaac tttcgaggca gatgggagag tacgtacaag agaggtttgt atggagaatt 2617
 gagtttgtg acggctagca ggctatgata tatgtageta tagtctagtc taaatctgaa 2677
 agaagagaac aagatgggtt gtccgaagag attgagagat caagcccggg catctgatgt 2737
 50 cgaacaaaca tgccctgggc tgccaacagt ttctagcaca ttatgaccat gttcatgtgt 2797
 aaattgggaa atgagccaga aaggtttatt atctaattca ttgattcatg cgactatgga 2857
 tacatatggg atttccagaa caaacagatg caacaaagca cggcatcttc caaagatcga 2917
 55 gtctcccccac aagtatgcgg caaggtttgt tgtaagaga tataaaagca gacgacaaaa 2977

EP 1 111 067 A2

caaatcgttt atcgaccctg tgcaccaaca ccgtgaccgt ttgacgagtt ggtagagttg 3037
 tagttgttgc tgttcaaagg agctccagac tggacgcttc caagcttcaa caacttctcg 3097
 5 gcagcgtcgc tgttcgggaa aagaaaaggc aaaaaggaac agagcgataa gcatatgtga 3157
 ttctctactt cttataggct cttagctcaa gtcaactcac atgtctttgg cggtagcgaa 3217
 gacgttctca agctgctgct tgggaagcttt tccgagcttg ccagtaggtc cctgggttga 3277
 10 gaagaagatg tcgaaggcta agggcgatga aaagcatgaa gatattagct atcggcgcgga 3337
 taaaagtgtg acgagatgaa aatggagaaa agatgattcg caccatcgac gacctcgacc 3397
 aaaggaatgg aggtgtcacc ggccttcac ttctgtact cctcaacgtt gacgaagatg 3457
 15 acgaagcagt cgggtggcctt agcctcgggc tcgtagatgc tgatgaaaca caataggtag 3517
 taggagagga gaaagagaag atgatgagat gtcaggatgc ttgcttcact gtagatggag 3577
 gaagaagata tgcgaagcaa gacatacact ttggaaagag cttgaaccat tgtag 3632
 20
 <210> 2
 <211> 3375
 <212> DNA
 <213> Phaffia rhodozyma
 25
 <220>
 <221> 5'UTR
 <222> (974)..(975)
 <223> EXPERIMENTAL
 30
 <220>
 <221> exon
 <222> (1040)..(1063)
 <220>
 <221> intron
 35 <222> (1064)..(1237)
 <220>
 <221> exon
 <222> (1238)..(1280)
 40
 <220>
 <221> intron
 <222> (1281)..(1358)
 <220>
 <221> exon
 45 <222> (1359)..(1402)
 <220>
 <221> intron
 <222> (1403)..(1483)
 50
 <220>
 <221> exon
 <222> (1484)..(1589)
 <220>
 <221> intron
 55 <222> (1590)..(1674)

EP 1 111 067 A2

```

<220>
<221> exon
<222> (1675)..(1826)

5   <220>
    <221> intron
    <222> (1827)..(1907)

    <220>
    <221> exon
10  <222> (1908)..(1922)

    <220>
    <221> intron
    <222> (1923)..(1992)

15  <220>
    <221> exon
    <222> (1993)..(2056)

    <220>
    <221> intron
20  <222> (2057)..(2131)

    <220>
    <221> exon
    <222> (2132)..(2157)

25  <220>
    <221> intron
    <222> (2158)..(2238)

    <220>
    <221> exon
30  <222> (2239)..(2293)

    <220>
    <221> intron
    <222> (2294)..(2376)

35  <220>
    <221> exon
    <222> (2377)..(2426)

    <220>
    <221> intron
40  <222> (2427)..(2524)

    <220>
    <221> exon
    <222> (2525)..(2542)

45  <220>
    <221> polyA_site
    <222> (2667)..(2668)
    <223> EXPERIMENTAL

50  <400> 2
    cttatccttc tgccgctggt ctctgtctgt cgagtgtgtg ggatgggttt ggatacgttc 60
    ctatacgaaa ggtagcgcag agcaaagctg acagtattaa gcaagacaag agcttctttc 120
    tgttgacaga tgaaaggacg aactatgaag ctgtccatgc tccccaaacc gattgacaca 180
55  ccgccgtcag gcaacgcaga atttctcact gcttcgacgt cacaccaaca tcgacccctc 240

```

EP 1 111 067 A2

alacctaaaa gcagatcgag acacattggt ggtcgccatg ttggatggat gtacatcaaa 300
 cccacagcat atatcactca catgtgagaa ctccgtagcc tctaccttct tgtctctcaa 360
 5 tctgaatgtc tcgttgagag gtggaatgaa tgtttacagt ttgagaagac gaaagaaaga 420
 aagagaagag aagagaggaa tacgtacgac gaagttatca tegtatggga acitttctaa 480
 aaaactgcct atagtagaga cgatctctgg aggaaagctc tgtagtatga tagtgaagag 540
 10 cgagcaagtc tgggcaagtg catccttcgt ctacaagaaa gagaccagga aatgaaggag 600
 agacgagtaa gcaggtacct accgatatgt gatcgttctc tctaccagc gatgccttca 660
 ccacgcgttc tatctcttct tgggatggca gatacatact taacgagagc aatctgatgt 720
 15 ataccgaact tcgaacggaa tgatcccaga atcctcttga acccttgaac ccttgaaccc 780
 tggaaaccaag taccaaccga gcaacacgcc gatacgggcc acaccacaga accacacgcc 840
 ctcgtcatta aaggtgggac gcgcgatgc tggttacgtt cggtcccaatc cggaagttac 900
 20 cggcttggac gtgcctgaa ccatgccctg acggtatttc gccttcagct aactccatct 960
 catcttttct ctttactacc acaaccacc cttgaacctt ctccccggc tttttacta 1020
 tatecatcta tcaatcacc atg gct cct tac act ctt ccc gac gtaagcttaa 1073
 25 agtttgagct gtgtgtgttt atctcaatct tggagtggaa ctcaacgttt ttgttttttg 1133
 ctctccgggt tttttatcgg catccctctt tttttccccc tcgtggctgc atatgatttg 1193
 ctcatcaatc ggcgtttccc atgcatcttt gtcacgtgtt tcag ctt cct tac gct 1249
 30 tac gat gcc ttg gag cct tac atc tct aag g gtgagattct tagtcagact 1300
 gttgttccgg ttcgacacga tagctaactg tctctcgttc ctcaatatga acatgcag 1358
 aa atc atg atc ctt cac cac tcc aag cac cat cag act tac gtc 1402
 35 gtacgtaate taaaggtcat ctccgtctac atggccggat caacttgctc atagatcttc 1462
 cttctgttcg gcgtacgta g acc aac ctg aac gcc gct atc cag gct ttc 1513
 tcc cag acc aat gac atc aag gcc cag atc gct ctt cag agc gct ctg 1561
 40 aag ttc aac gga gga gga cac atc aac c gtacgatcat tctccctctt 1609
 ctggcttate atatgtgttg ctgtcacta acacgcacgc aaccccgga tatctcacc 1669
 tgtag ac tcc ctg ttc tgg aag aac atg gct cct gcc gac tct gct gat 1718
 45 gcc aag ctg acc gag gga tgc ctg aag act gcc atc gac aag gac ttt 1766
 gga tcc ttc gag gag ttc aag aag aag ttc aac act gct act ctg ggt 1814
 gtc cag gga tct gtcagtatct cgttctcttc gacatactct cagctttcct 1866
 50 tccgtaaaact gacgaatagt ttttcggaca tgtacttgta g gga tgg gga tgg ctg 1922
 gtgcgtttga cctttttcca ctttgaaca tagcgatagt gatacctaac aactgtgaat 1982
 tggaatatag gga tac aac acc gct acc aag cac ctg gag atc gcc acc 2031
 55 acc gcc aac cag gat ccc ctt atc a gtatgtgact tctctcgtgt 2076

EP 1 111 067 A2

5 ggccaccata agccagttgc tgacacattc cgctcgtctg ctctcgactt cglag ct 2133
 ttg act ccc acc att ggt ctt gac gttagtaatt ctatctagtg attggagtcg 2187
 agttctgaac ttgccttgat ctcaaacgaa tgaatcaatt tcttttggtg g atc tgg 2244
 gag cac gct ttc tac ctg cag tac aag aat gtc aag cct gat tac ctt g 2293
 gtacgtaatt ctctattcgt ttgccccggg ttgatctttg actcactctt caaaatgttc 2353
 10 tcttttgta actttgaaaa acag cc gct ttc tgg aac gtc tgc aac ttt gct 2405
 gag gct cag cga agg ttt gat gtgagtagag ggcctacccc tacggaggaa 2456
 gcgaagggtga gctgaccact ttttatcttc ctgatttggg atgaacgac cgatgatcaa 2516
 15 acaaacag gct gct gtc aag gct taa tggctccatt tatctctttg attcgacggc 2572
 gatgacggct ttctcgctac cgaagaaggc aaggctatga ttactgttat tctgccatgt 2632
 tgcttgcttt gctatgctct atgttctttt cttttgcctc tcttcaaagc caaggcggtta 2692
 20 aggaaggccc ttcagtctgt ttacatattg cacatataca tgagaacata tcacggactc 2752
 ggcggtctgt ggtctctctg agcgtcggc tcaagattag tgtccacacg tgaagcgttc 2812
 ggtgccatcc aacctggtag gaatcccca cgggcgggaa tccaattatc aattggcggg 2872
 25 cggccagatt cgagctcggg tatctcagaa gcgtcaagcg ggcgcatttc caggccttta 2932
 agaggagcaa atttaatcgg cctgggtgtt cagcgagaca cgaacagttt gaaacagagt 2992
 ctgcttggtga gttactcggc gagatcactg aggactaaac tttctcagct cgtggacgaa 3052
 30 aagaacgaac caaacggctt tccctgtatc tcgaccatct ccttctccat ctcttacaac 3112
 acctcggatg aactccaagg cttgcttttc aaagttcaaa caaactcagg gttgccatcc 3172
 acctggtttg tctctaacga gccgagggat atccatcgtt cggaacgttt gaacagactg 3232
 35 gatggttaggt ggccggctcg ttcggaagcc aatcataatg gtgggaatcg agagaaggaa 3292
 tgattgggcc cagtgtttta gacttgtgtt tgttggcaga gtacggacgg aaagtaggac 3352
 agacttaatc aaggcgagcc aag 3375

40
 <210> 3
 <211> 951
 <212> DNA
 <213> *Phaffia rhodozyma*

45
 <220>
 <221> exon
 <222> (1)..(34)

50
 <220>
 <221> intron
 <222> (35)..(115)

<220>
 <221> exon
 <222> (117)..(159)

55
 <220>
 <221> intron

EP 1 111 067 A2

<222> (160)..(242)

<220>
<221> exon
5 <222> (243)..(363)

<220>
<221> intron
<222> (364)..(436)

10 <220>
<221> exon
<222> (437)..(951)

<400> 3

15 tcc gga agc tca gat acc gct cga gat cct cga g gtttgtgtgc 44

tttcgctttg ttcgcatgga tgaagctggt aacttaaaaa aatcctcgtg tttctctttg 104

tttcaacata g qt ttc tct ctt aag gtc aag acc tct gag gga aac tgg 153

20 gac ttt gtacgtattc ttatcgactg agtcatcaag ctctgtatcg ctctcttacc 209

ctcatccttt tgtgtctctg tctacacctc tag gtc gga aac aac act ccc atc 263

ttt ttc ttg aga gac cca gcc aag ttt ccg atc ttc att cac acc cag 311

25 aag agg aac ccg cag aca aac tct aaa gac aag gac gct ttc tgg gac 359

tac c gtctgtata accttgtcac tccctgcgtg ccgtctctgat tcatgttgac 412

cttgtctttg atataatttt atag ta tcc caa aac ccc gag tcc gtg cat cag 465

30 gtc ctg cac ctg ttc agt gat cga gga acc cct gct tct tac cga cac 513

atg cat ggt tac tct gga cac acc ttc aag atg gtc aac agg aac ggt 561

gac tgg aat tat gtc cag att cac atg cgc acc gat cag ggt gtc aag 609

35 act cac acc aat gaa gag gct tgc aaa ctg gac gcc tcc aat ccc gat 657

tca aac gga gac gac ttg ttc gac gca atc aag aat gga gac ttc cct 705

agc tgg acg gtt cag gta cag gta atg tct cct gag cag gcc cag aag 753

40 ttc aga tac aac att ctg gat ctg acc aag gtc tgg tcc cac aag gag 801

ttc cca ctt agg acg att gga aag ttc act ttg aac cga aac gtg gat 849

aac tat ttc gca gag gtc gaa cag ctg gcc ttt gct cct tcc cat ctg 897

45 cct cct gga atc gag ccc tgc aac gat ccc gtc ctt cag gct cga cta 945

ttc tcc 951

<210> 4
50 <211> 669
<212> DNA
<213> Phaffia rhodozyma

<220>
<221> CDS
55 <222> (1)..(669)

<400> 4

EP 1 111 067 A2

	atg tct gtt cga gca tcc ctc tct tcc gtg tct aga cag act ttc gtc	48
	Met Ser Val Arg Ala Ser Leu Ser Ser Val Ser Arg Gln Thr Phe Val	
	1 5 10 15	
5	gct cct gct gct ttc cag atc agg gca aag cat acc ctg cct gag ctt	96
	Ala Pro Ala Ala Phe Gln Ile Arg Ala Lys His Thr Leu Pro Glu Leu	
	20 25 30	
10	cct tac gct tac gat gcc ctg gag ccc tcc atc tcc aag gag atc atg	144
	Pro Tyr Ala Tyr Asp Ala Leu Glu Pro Ser Ile Ser Lys Glu Ile Met	
	35 40 45	
15	acc ctt cac cac acc aag cac cat cag act tat gtt aac ggc ctc aac	192
	Thr Leu His His Thr Lys His His Gln Thr Tyr Val Asn Gly Leu Asn	
	50 55 60	
20	gct gcc gag gag agc tac tcg gcc gct gtg ggc aag gag gat gtg ctt	240
	Ala Ala Glu Glu Ser Tyr Ser Ala Ala Val Gly Lys Glu Asp Val Leu	
	65 70 75 80	
25	acc cag gtt aag ctt cag tct gct ctc aag ttc aac gga gga gga cac	288
	Thr Gln Val Lys Leu Gln Ser Ala Leu Lys Phe Asn Gly Gly Gly His	
	85 90 95	
30	atc aat cac tct ctg ttc tgg aag aac ttg gct ccc tat gga tcc gag	336
	Ile Asn His Ser Leu Phe Trp Lys Asn Leu Ala Pro Tyr Gly Ser Glu	
	100 105 110	
35	gag gct acc ctc tct gaa gga cct ctc aag aag gct atc gag gaa tct	384
	Glu Ala Thr Leu Ser Glu Gly Pro Leu Lys Lys Ala Ile Glu Glu Ser	
	115 120 125	
40	ttt ggt tct ttc gag gcc ttc aag aag aag ttc aac gct gac acc gct	432
	Phe Gly Ser Phe Glu Ala Phe Lys Lys Lys Phe Asn Ala Asp Thr Ala	
	130 135 140	
45	gct gtc caa gga tcc gga tgg ggc tgg ctt ggc ttg aac ccg ctt act	480
	Ala Val Gln Gly Ser Gly Trp Gly Trp Leu Gly Leu Asn Pro Leu Thr	
	145 150 155 160	
50	aag aag ctg gaa gtc acc acg acc gcc aac cag gac cct ctg ctt act	528
	Lys Lys Leu Glu Val Thr Thr Thr Ala Asn Gln Asp Pro Leu Leu Thr	
	165 170 175	
55	cac att cct atc atc gga gtt gac atc tgg gag cac gct ttc tac ctt	576
	His Ile Pro Ile Ile Gly Val Asp Ile Trp Glu His Ala Phe Tyr Leu	
	180 185 190	
60	cag tac aag aac gtc aag cct gac tat ctc gct gct gtt tgg tcc gtt	624
	Gln Tyr Lys Asn Val Lys Pro Asp Tyr Leu Ala Ala Val Trp Ser Val	
	195 200 205	
65	atc aac tac aag gag gca gag gcc cga ttg cag gct gct ctc taa	669
	Ile Asn Tyr Lys Glu Ala Glu Ala Arg Leu Gln Ala Ala Leu	
	210 215 220	
70	<210> 5	
	<211> 222	
	<212> PRT	
	<213> Phaffia rhodozyma	
75	<400> 5	
	Met Ser Val Arg Ala Ser Leu Ser Ser Val Ser Arg Gln Thr Phe Val	
	1 5 10 15	

EP 1 111 067 A2

Ala Pro Ala Ala Phe Gln Ile Arg Ala Lys His Thr Leu Pro Glu Leu
20 25 30

5 Pro Tyr Ala Tyr Asp Ala Leu Glu Pro Ser Ile Ser Lys Glu Ile Met
35 40 45

Thr Leu His His Thr Lys His His Gln Thr Tyr Val Asn Gly Leu Asn
50 55 60

10 Ala Ala Glu Glu Ser Tyr Ser Ala Ala Val Gly Lys Glu Asp Val Leu
65 70 75 80

Thr Gln Val Lys Leu Gln Ser Ala Leu Lys Phe Asn Gly Gly Gly His
85 90 95

15 Ile Asn His Ser Leu Phe Trp Lys Asn Leu Ala Pro Tyr Gly Ser Glu
100 105 110

Glu Ala Thr Leu Ser Glu Gly Pro Leu Lys Lys Ala Ile Glu Glu Ser
115 120 125

20 Phe Gly Ser Phe Glu Ala Phe Lys Lys Lys Phe Asn Ala Asp Thr Ala
130 135 140

Ala Val Gln Gly Ser Gly Trp Gly Trp Leu Gly Leu Asn Pro Leu Thr
145 150 155 160

25 Lys Lys Leu Glu Val Thr Thr Thr Ala Asn Gln Asp Pro Leu Leu Thr
165 170 175

His Ile Pro Ile Ile Gly Val Asp Ile Trp Glu His Ala Phe Tyr Leu
180 185 190

30 Gln Tyr Lys Asn Val Lys Pro Asp Tyr Leu Ala Ala Val Trp Ser Val
195 200 205

Ile Asn Tyr Lys Glu Ala Glu Ala Arg Leu Gln Ala Ala Leu
210 215 220

35

<210> 6
<211> 597
<212> DNA
<213> Phaffia rhodozyma

40

<220>
<221> CDS
<222> (1)..(597)

<400> 6

45 atg gct cct tac act ctt ccc gac ctt cct tac gct tac gat gcc ttg 48
Met Ala Pro Tyr Thr Leu Pro Asp Leu Pro Tyr Ala Tyr Asp Ala Leu
1 5 10 15

gag cct tac atc tct aag gaa atc atg atc ctt cac cac tcc aag cac 96
Glu Pro Tyr Ile Ser Lys Glu Ile Met Ile Leu His His Ser Lys His
20 25 30

50 cat cag act tac gtc acc aac ctc aac gcc gct atc cag gct ttc tcc 144
His Gln Thr Tyr Val Thr Asn Leu Asn Ala Ala Ile Gln Ala Phe Ser
35 40 45

55 cag acc aat gac atc aag gcc cag atc gct ctt cag agc gct ctc aag 192
Gln Thr Asn Asp Ile Lys Ala Gln Ile Ala Leu Gln Ser Ala Leu Lys
50 55 60

EP 1 111 067 A2

ttc aac gga gga gga cac atc aac cac tcc ctc ttc tgg aag aac atg 240
 Phe Asn Gly Gly Gly His Ile Asn His Ser Leu Phe Trp Lys Asn Met
 65 70 75 80
 5 gct cct gcc gac tct gct gat gcc aag ctc acc gag gga tgg ctc aag 288
 Ala Pro Ala Asp Ser Ala Asp Ala Lys Leu Thr Glu Gly Ser Leu Lys
 85 90 95
 10 act gcc atc gac aag gac ttt gga tcc ttc gag gag ttc aag aag aag 336
 Thr Ala Ile Asp Lys Asp Phe Gly Ser Phe Glu Glu Phe Lys Lys Lys
 100 105 110
 ttc aac act gct act ctc ggt gtc cag gga tct gga tgg gga tgg ctc 384
 Phe Asn Thr Ala Thr Leu Gly Val Gln Gly Ser Gly Trp Gly Trp Leu
 115 120 125
 15 gga tac aac acc gct acc aag cac ctc gag atc gcc acc acc gcc aac 432
 Gly Tyr Asn Thr Ala Thr Lys His Leu Glu Ile Ala Thr Thr Ala Asn
 130 135 140
 20 cag gat ccc ctt atc act ttg act ccc atc att ggt ctt gac atc tgg 480
 Gln Asp Pro Leu Ile Thr Leu Thr Pro Ile Ile Gly Leu Asp Ile Trp
 145 150 155 160
 gag cac gct ttc tac ctc cag tac aag aat gtc aag cct gat tac ctt 528
 Glu His Ala Phe Tyr Leu Gln Tyr Lys Asn Val Lys Pro Asp Tyr Leu
 165 170 175
 25 gcc gct ttc tgg aac gtc tgc aac ttt gct gag gct cag cga agg ttt 576
 Ala Ala Phe Trp Asn Val Cys Asn Phe Ala Glu Ala Gln Arg Arg Phe
 180 185 190
 30 gat gct gct gtc aag gct taa 597
 Asp Ala Ala Val Lys Ala
 195
 35 <210> 7
 <211> 198
 <212> PRT
 <213> Phaffia rhodozyma
 <400> 7
 Met Ala Pro Tyr Thr Leu Pro Asp Leu Pro Tyr Ala Tyr Asp Ala Leu
 1 5 10 15
 40 Glu Pro Tyr Ile Ser Lys Glu Ile Met Ile Leu His His Ser Lys His
 20 25 30
 His Gln Thr Tyr Val Thr Asn Leu Asn Ala Ala Ile Gln Ala Phe Ser
 35 40 45
 45 Gln Thr Asn Asp Ile Lys Ala Gln Ile Ala Leu Gln Ser Ala Leu Lys
 50 55 60
 Phe Asn Gly Gly Gly His Ile Asn His Ser Leu Phe Trp Lys Asn Met
 65 70 75 80
 50 Ala Pro Ala Asp Ser Ala Asp Ala Lys Leu Thr Glu Gly Ser Leu Lys
 85 90 95
 Thr Ala Ile Asp Lys Asp Phe Gly Ser Phe Glu Glu Phe Lys Lys Lys
 100 105 110
 55 Phe Asn Thr Ala Thr Leu Gly Val Gln Gly Ser Gly Trp Gly Trp Leu

EP 1 111 067 A2

	115	120	125	
	Gly Tyr Asn Thr Ala Thr Lys His Leu Glu Ile Ala Thr Thr Ala Asn			
	130	135	140	
5	Gln Asp Pro Leu Ile Thr Leu Thr Pro Ile Ile Gly Leu Asp Ile Trp			
	145	150	155	160
	Glu His Ala Phe Tyr Leu Gln Tyr Lys Asn Val Lys Pro Asp Tyr Leu			
		165	170	175
10	Ala Ala Phe Trp Asn Val Cys Asn Phe Ala Glu Ala Gln Arg Arg Phe			
		180	185	190
	Asp Ala Ala Val Lys Ala			
	195			
15				
	<210> 8			
	<211> 714			
	<212> DNA			
20	<213> Phaffia rhodozyma			
	<220>			
	<221> CDS			
	<222> (1)..(714)			
25	<400> 8			
	tcc gga agc tca gat acc gct cga gat cct cga ggt ttc tct ctt aag			48
	Ser Gly Ser Ser Asp Thr Ala Arg Asp Pro Arg Gly Phe Ser Leu Lys			
	1	5	10	15
	gtc aag acc tct gag gga aac tgg gac ttt gtc gga aac aac act ccc			96
30	Val Lys Thr Ser Glu Gly Asn Trp Asp Phe Val Gly Asn Asn Thr Pro			
		20	25	30
	atc ttt ttc ttg aga gac cca gcc aag ttt ccg atc ttc att cac acc			144
	Ile Phe Phe Leu Arg Asp Pro Ala Lys Phe Pro Ile Phe Ile His Thr			
		35	40	45
35	cag aag agg aac ccg cag aca aac tct aaa gac aag gac gct ttc tgg			192
	Gln Lys Arg Asn Pro Gln Thr Asn Ser Lys Asp Lys Asp Ala Phe Trp			
		50	55	60
	gac tac cta tcc caa aac ccc gag tcc gtg cat cag gtg ctg cac ctg			240
40	Asp Tyr Leu Ser Gln Asn Pro Glu Ser Val His Gln Val Leu His Leu			
		65	70	75
	ttc agt gat cga gga acc cct gct tct tac cga cac atg cat ggt tac			288
	Phe Ser Asp Arg Gly Thr Pro Ala Ser Tyr Arg His Met His Gly Tyr			
		85	90	95
45	tct gga cac acc ttc aag atg gtc aac agg aac ggt gac tgg aat tat			336
	Ser Gly His Thr Phe Lys Met Val Asn Arg Asn Gly Asp Trp Asn Tyr			
		100	105	110
	gtc cag att cac atg cgc acc gat cag ggt gtc aag act cac acc aat			384
50	Val Gln Ile His Met Arg Thr Asp Gln Gly Val Lys Thr His Thr Asn			
		115	120	125
	gaa gag gct tcg aaa ctc gac gcc tcc aat ccc gat tca aac gga gac			432
	Glu Glu Ala Ser Lys Leu Asp Ala Ser Asn Pro Asp Ser Asn Gly Asp			
		130	135	140
55	gac ttg ttc gac gca atc aag aat gga gac ttc cct agc tgg acg gtt			480

EP 1 111 067 A2

Asp Leu Phe Asp Ala Ile Lys Asn Gly Asp Phe Pro Ser Trp Thr Val
145 150 155 160

5 cag gta cag gta atg tct cct gag cag gcc cag aag ttc aga tac aac 528
Gln Val Gln Val Met Ser Pro Glu Gln Ala Gln Lys Phe Arg Tyr Asn
165 170 175

att ctg gat ctg acc aag gtc tgg tcc cac aag gag ttc cca ctt agg 576
Ile Leu Asp Leu Thr Lys Val Trp Ser His Lys Glu Phe Pro Leu Arg
180 185 190

10 acg att gga aag ttc act ttg aac cga aac gtg gat aac tat ttc gca 624
Thr Ile Gly Lys Phe Thr Leu Asn Arg Asn Val Asp Asn Tyr Phe Ala
195 200 205

gag gtt gaa cag ctg gcc ttt gct cct tcc cat ctg cct cct gga atc 672
Glu Val Glu Gln Leu Ala Phe Ala Pro Ser His Leu Pro Pro Gly Ile
210 215 220

15 gag ccc tcg aac gat ccc gtc ctt cag gct cga cta ttc tcc 714
Glu Pro Ser Asn Asp Pro Val Leu Gln Ala Arg Leu Phe Ser
225 230 235

20 <216> 9
<211> 238
<212> PRT
<213> Phaffia rhodozyma

25 <400> 9
Ser Gly Ser Ser Asp Thr Ala Arg Asp Pro Arg Gly Phe Ser Leu Lys
1 5 10 15

30 Val Lys Thr Ser Glu Gly Asn Trp Asp Phe Val Gly Asn Asn Thr Pro
20 25 30

Ile Phe Phe Leu Arg Asp Pro Ala Lys Phe Pro Ile Phe Ile His Thr
35 40 45

Gln Lys Arg Asn Pro Gln Thr Asn Ser Lys Asp Lys Asp Ala Phe Trp
50 55 60

35 Asp Tyr Leu Ser Gln Asn Pro Glu Ser Val His Gln Val Leu His Leu
65 70 75 80

40 Phe Ser Asp Arg Gly Thr Pro Ala Ser Tyr Arg His Met His Gly Tyr
85 90 95

Ser Gly His Thr Phe Lys Met Val Asn Arg Asn Gly Asp Trp Asn Tyr
100 105 110

45 Val Gln Ile His Met Arg Thr Asp Gln Gly Val Lys Thr His Thr Asn
115 120 125

Glu Glu Ala Ser Lys Leu Asp Ala Ser Asn Pro Asp Ser Asn Gly Asp
130 135 140

50 Asp Leu Phe Asp Ala Ile Lys Asn Gly Asp Phe Pro Ser Trp Thr Val
145 150 155 160

Gln Val Gln Val Met Ser Pro Glu Gln Ala Gln Lys Phe Arg Tyr Asn
165 170 175

55 Ile Leu Asp Leu Thr Lys Val Trp Ser His Lys Glu Phe Pro Leu Arg
180 185 190

EP 1 111 067 A2

Thr Ile Gly Lys Phe Thr Leu Asn Arg Asn Val Asp Asn Tyr Phe Ala
195 200 205

5 Glu Val Glu Gln Leu Ala Phe Ala Pro Ser His Leu Pro Pro Gly Ile
210 215 220

Glu Pro Ser Asn Asp Pro Val Leu Gln Ala Arg Leu Phe Ser
225 230 235

10 <210> 10
<211> 23
<212> DNA
<213> Artificial Sequence

15 <220>
<223> Description of Artificial Sequence:Sod1(sense
primer for cloning of SOD genes)

<400> 10
aarcaycayc aractaygt naa 23

20 <210> 11
<211> 23
<212> DNA
<213> Artificial Sequence

25 <220>
<223> Description of Artificial Sequence:Sod4 (antisense
primer for cloning of SOD genes)

<400> 11
gcccancng anccytgnac ncc 23

30 <210> 12
<211> 26
<212> DNA
<213> Artificial Sequence

35 <220>
<223> Description of Artificial Sequence:Sod14 (sense
primer for the construction of SOD1-disrupting
plasmid)

40 <400> 12
ggtacctcgg atgataggaa tgtgag 26

45 <210> 13
<211> 26
<212> DNA
<213> Artificial Sequence

50 <220>
<223> Description of Artificial Sequence:Sod15
(antisense primer for the construction of
SOD1-disrupting plasmid)

<400> 13
gaattcagtt caacggagga ggacac 26

55 <210> 14
<211> 26

<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Sod47 (sense
primer for the construction of SOD2-disrupting
plasmid)

<400> 14
gaattcggag gaggacacat caaccg

26

<210> 15
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Sod48
(antisense primer for the construction of
SOD2-disrupting plasmid)

<400> 15
ggtacctgta ctggaggtag aaagcg

26

<210> 16
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Sod2 (sense
primer for cloning of CAT gene)

<400> 16
mgnttytcna cngtnggngg nga

23

<210> 17
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Cat5 (antisense
primer for cloning of CAT gene)

<400> 17
ckrtgackyt gngtrtcngg rta

23

Claims

1. A process for producing carotenoids which comprises cultivating a recombinant organism whose gene for one or more active oxygen species-quenching factor(s) is substantially disrupted with the aid of a disruption cassette specific to the said gene, and recovering carotenoids from the culture.
2. A process according to claim 1, wherein the host organism of the said recombinant organism belongs to the kingdom of *Monera*, *Protista* or *Fungi*.

3. A process according to claim 1 or 2, wherein the host organism of the said recombinant organism belongs to the genus *Erwinia*, *Rhodobacter*, *Myxococcus*, *Flavobacter*, *Paracoccus*, *Synechococcus*, *Synechocystis*, *Agrobacterium*, *Streptomyces*, *Haematococcus*, *Dunaliella*, *Phaffia*, *Xanthophyllomyces*, *Neurospora*, *Rhodotorula*, *Blakeslea*, or *Phycomyces*.

4. A process according to claim 3, wherein the host organism of the said recombinant organism is a strain of *P. rhodozyma*

5. A process according to any one of claims 1 - 4, wherein the active oxygen species-quenching factor(s) is(are) mitochondrial superoxide dismutase (SOD), cytoplasmic superoxide dismutase (SOD) and/or catalase.

6. A recombinant organism producible of carotenoids, which is characterized in that the gene for at least one active oxygen species-quenching factor is substantially disrupted with the aid of introduction of a disruption cassette specific to the said gene.

7. A recombinant organism according to claim 6, wherein the host organism of the said recombinant organism belongs to the kingdom of *Monera*, *Protista* or *Fungi*, more preferably to the genus *Erwinia*, *Rhodobacter*, *Myxococcus*, *Flavobacter*, *Paracoccus*, *Synechococcus*, *Synechocystis*, *Agrobacterium*, *Streptomyces*, *Haematococcus*, *Dunaliella*, *Phaffia*, *Xanthophyllomyces*, *Neurospora*, *Rhodotorula*, *Blakeslea*, or *Phycomyces*.

8. A recombinant organism according to claim 6 or 7, wherein the active oxygen species-quenching factor(s) to be disrupted is(are) mitochondrial superoxide dismutase (SOD), cytoplasmic superoxide dismutase (SOD) and/or catalase.

9. A disruption cassette for disrupting a gene coding for an active oxygen species-quenching factor effective on carotenogenesis in an organism which is producible of carotenoids, which comprises a partial nucleotide sequence substantially identical to a part of the DNA sequence coding for an active oxygen species-quenching factor and a selectable marker gene.

10. A disruption cassette according to claim 9, wherein the said organism belongs to the kingdom of *Monera*, *Protista* or *Fungi*, more preferably to the genus *Erwinia*, *Rhodobacter*, *Myxococcus*, *Flavobacter*, *Paracoccus*, *Synechococcus*, *Synechocystis*, *Agrobacterium*, *Streptomyces*, *Haematococcus*, *Dunaliella*, *Phaffia*, *Xanthophyllomyces*, *Neurospora*, *Rhodotorula*, *Blakeslea*, or *Phycomyces*.

11. A disruption cassette according to claim 9 or 10, wherein the active oxygen species-quenching factor(s) to be disrupted is(are) mitochondrial superoxide dismutase (SOD), cytoplasmic superoxide dismutase (SOD) and/or catalase.

12. A disruption cassette according to any one of claims 9 to 11, wherein the partial nucleotide sequence coding for an active oxygen species-quenching factor is identical to at least a part of the DNA sequence coding for the active oxygen species-quenching factor of the organism into which the disruption cassette is to be introduced.

13. A disruption cassette according to claim 12, wherein the partial nucleotide sequence substantially identical to a part of the DNA sequence coding for an active oxygen species-quenching factor comprises at least one deletion and/or mutation compared to the corresponding functional gene.

14. A recombinant DNA sequence coding for an active oxygen species-quenching factor effective on carotenogenesis in an organism producible of carotenoids.

15. A recombinant DNA sequence according to claim 14, wherein the said organism belongs to the kingdom of *Monera*, *Protista* or *Fungi*.

16. A recombinant DNA sequence according to claim 14 or 15, wherein the said organism belongs to the kingdom of *Monera*, *Protista* or *Fungi*, more preferably to the genus *Erwinia*, *Rhodobacter*, *Myxococcus*, *Flavobacter*, *Paracoccus*, *Synechococcus*, *Synechocystis*, *Agrobacterium*, *Streptomyces*, *Haematococcus*, *Dunaliella*, *Phaffia*, *Xanthophyllomyces*, *Neurospora*, *Rhodotorula*, *Blakeslea*, or *Phycomyces*.

17. A recombinant DNA sequence according to any one of claims 14 - 16, which is a sequence originated from a

microorganism of *P. rhodozyma*.

18. A recombinant DNA sequence according to any one of claims 14 - 17, wherein the active oxygen species-quenching factor is mitochondrial superoxide dismutase.

19. A recombinant DNA sequence according to claim 18, wherein the sequence is identified by SEQ ID NO: 1 or 4, or has high homology to the sequence of SEQ ID NO: 1 or 4 enough to recombine with either of the sequences SEQ ID NOs: 1 and 4.

20. A recombinant DNA sequence according to claim 14 - 17, wherein the active oxygen species-quenching factor is cytoplasmic superoxide dismutase.

21. A recombinant DNA sequence according to claim 20, wherein the sequence is identified by SEQ ID NO: 2 or 6, or has high homology to the sequence of SEQ ID NO: 2 or 6 enough to recombine with either of the sequences of SEQ ID NOs: 2 and 6.

22. A recombinant DNA sequence according to claim 14 - 17, wherein the active oxygen species-quenching factor is catalase.

23. A recombinant DNA sequence according to claim 22, wherein the sequence is identified by SEQ ID NO: 3 or 8, or has high homology to the sequence of SEQ ID NO: 3 or 8 enough to recombine with either of the sequences of SEQ ID NOs: 3 and 8.

24. A recombinant DNA fragment which comprises a coding region for a transit peptide upstream the coding region of an objective protein.

25. A recombinant DNA fragment according to claim 24, wherein the objective protein is mitochondrial superoxide dismutase.

26. A method for locating an objective protein in mitochondria which comprises expressing the recombinant DNA fragment defined in claim 24 or 25 in an appropriate recombinant host organism.

27. A polynucleotide for use as probe or primer for cloning gene for the active oxygen species-quenching factor(s) effective on carotenogenesis in an organism producible of carotenoids.

28. A polynucleotide according to claim 27, wherein the active oxygen species-quenching factor(s) is(are) mitochondrial superoxide dismutase (SOD), cytoplasmic superoxide dismutase (SOD) and/or catalase.





FIG. 1 Activity staining for super oxide dismutase in SOD disruptants and their host strain, ATCC96594
 lane 1 & 4 ; ATCC96594
 lane 2 ; ATCC96594 :: pSOD/G717 (Δ SOD1)
 lane 3 ; ATCC96594 :: pSOD/G828 (Δ SOD2)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 111 067 A3**

(12) **EUROPEAN PATENT APPLICATION**

(88) Date of publication A3:
10.10.2001 Bulletin 2001/41

(51) Int Cl.7: **C12P 23/00**, C12N 9/08,
C12N 9/02

(43) Date of publication A2:
27.06.2001 Bulletin 2001/26

(21) Application number: **00126114.8**

(22) Date of filing: **29.11.2000**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **01.12.1999 EP 99123821**

(71) Applicant: **F. HOFFMANN-LA ROCHE AG
4070 Basel (CH)**

(72) Inventors:
• **Hoshino, Tatsuo**
Kamakura-shi, Kanagawa-ken 248-0027 (JP)

• **Ojima, Kazuyuki**
Fujisawa-shi, Kanagawa-ken 251-0877 (JP)
• **Setoguchi, Yutaka**
Fujisawa-shi, Kanagawa-ken 251-0033 (JP)

(74) Representative: **Braun, Axel et al**
F.Hoffmann-La Roche AG
Patent Department (PLP),
124 Grenzacherstrasse
4070 Basel (CH)

(54) **Recombinant production of carotenoids, particularly astaxanthin**

(57) The present invention is directed to a process for producing carotenoids comprising cultivating a recombinant organism having a gene for one or more active oxygen species-quenching factor(s) that is substantially disrupted with a disruption cassette specific to the

gene, and recovering carotenoids from the culture, as well as to genetic materials useful in the said process, such as a recombinant organism producible of carotenoids, a disruption cassette, a recombinant DNA sequence, a recombinant DNA fragment; and a polynucleotide.

EP 1 111 067 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 00 12 6114

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
D,A	SCHROEDER WILLIAM A ET AL: "Antioxidant role of carotenoids in Phaffia rhodozyma." JOURNAL OF GENERAL MICROBIOLOGY, vol. 139, no. 5, 1993, pages 907-912, XP000993067 ISSN: 0022-1287 * the whole document *	1-13	C12P23/00 C12N9/08 C12N9/02
A	AN GIL-HWAN: "Photosensitization of the yeast Phaffia rhodozyma at a low temperature for screening carotenoid hyperproducing mutants." APPLIED BIOCHEMISTRY AND BIOTECHNOLOGY, vol. 66, no. 3, 1997, pages 263-268, XP000993071 ISSN: 0273-2289 * the whole document *	1-13	
D,A	SHAISH AVIV ET AL: "Are active oxygen species involved in induction of beta-carotene in Dunaliella bardawil?" PLANTA (HEIDELBERG), vol. 190, no. 3, 1993, pages 363-368, XP000993142 ISSN: 0032-0935 * the whole document *	1-13	TECHNICAL FIELDS SEARCHED (Int.Cl.7) C12P C12N
A	SCHROEDER WA ET AL.: "Singlet oxygen and peroxy radicals regulate carotenoid biosynthesis in Phaffia rhodozyma." J BIOL CHEM 1995 AUG 4;270(31):18374-9., XP000577110 * the whole document *	1-13	
-/--			
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 14 August 2001	Examiner Andres, S
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application I : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPC FORM 1503 03 82 (04/01)



European Patent
Office

Application Number

EP 00 12 6114

CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing more than ten claims.

- ☐ Only part of the claims have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid, namely claim(s):
- ☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.

LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

see sheet B

- ☒ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.
- ☐ As all searchable claims could be searched without effort justifying an additional fee, the Search Division did not invite payment of any additional fee.
- ☐ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid, namely claims:
- ☐ None of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims, namely claims:



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 00 12 6114

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A	MARTINEZ C ET AL.: "Genetic transformation of astaxanthin mutants of <i>Phaffia rhodozyma</i> ." ANTONIE VAN LEEUWENHOEK 1998 FEB:73(2):147-53., XP000993172		
X	CORTEZ NESTOR ET AL: "Molecular cloning and expression analysis of the <i>Rhodobacter capsulatus</i> <i>sodB</i> gene, encoding an iron superoxide dismutase." JOURNAL OF BACTERIOLOGY, vol. 180, no. 20, October 1998 (1998-10), pages 5413-5420, XP002166756 ISSN: 0021-9193 * the whole document *	6-16,20, 27,28	
X	JAKOPITSCH CHRISTA ET AL: "Catalase-peroxidase from the cyanobacterium <i>Synechocystis</i> PCC 6803: Cloning, overexpression in <i>Escherichia coli</i> , and kinetic characterization." BIOLOGICAL CHEMISTRY, vol. 380, no. 9, September 1999 (1999-09), pages 1087-1096, XP000993333 ISSN: 1431-6730 * the whole document *	14-16, 22,27,28	TECHNICAL FIELDS SEARCHED (Int.Cl.7)
X	FORKL HUBERT ET AL: "Molecular cloning, sequence analysis and expression of the gene for catalase-peroxidase (<i>cpeA</i>) from the photosynthetic bacterium <i>Rhodobacter capsulatus</i> B10." EUROPEAN JOURNAL OF BIOCHEMISTRY, vol. 214, no. 1, 1993, pages 251-258, XP000993284 ISSN: 0014-2956 * the whole document *	14-16, 22,27,28	
-/--			
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 14 August 2001	Examiner Andres, S
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document</p> <p>T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons & member of the same patent family, corresponding document</p>			

EPC FORM 1503 03 52 (F04C01)



European Patent
Office

**LACK OF UNITY OF INVENTION
SHEET B**

Application Number
EP 00 12 6114

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

1. Claims: 1-23,27-28

1.1. Claims: 1-13

A process for producing carotenoids in an organism by disrupting a gene encoding an oxygen species quenching factor, and the disruption cassette used.

1.2. Claims: 14-17,27-28 (all partially) and claims 18-19

A recombinant DNA coding for a mitochondrial SOD of a carotenoid producing organism.

1.3. Claims: 14-17,27-28 (all partially) and claims 20-21

A recombinant DNA coding for a cytoplasmic SOD of a carotenoid producing organism.

1.4. Claims: 14-17,27-28 (all partially) and claims 22-23

A recombinant DNA coding for a catalase of a carotenoid producing organism.

2. Claims: 24-26

A recombinant DNA comprising a coding region for a transit peptide upstream of the coding region for a protein, and its use for locating the protein in mitochondria.

Please note that all inventions mentioned under item 1, although not necessarily linked by a common inventive concept, could be searched without effort justifying an additional fee.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 00 12 6114

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	CHUNG H -J ET AL: "Duplicate genes for Fe-containing superoxide dismutase in Streptomyces coelicolor A3(2)" GENE,NL,ELSEVIER BIOMEDICAL PRESS. AMSTERDAM, vol. 231, no. 1-2, 29 April 1999 (1999-04-29), pages 87-93, XP004166781 ISSN: 0378-1119 * the whole document *	6-16,20, 27,28	
X	WO 90 02804 A (PLANT GENETIC SYSTEMS NV) 22 March 1990 (1990-03-22) * abstract * * page 7, line 19 - page 8, line 17 * * page 13, line 9 - page 14, line 22 * * claims *	24-26	
X	CREISSEN GARY ET AL: "Simultaneous targeting of pea glutathione reductase and of a bacterial fusion protein to chloroplasts and mitochondria in transgenic tobacco." PLANT JOURNAL, vol. 8, no. 2, 1995, pages 167-175, XP002174824 ISSN: 0960-7412 * pGR40 *	24-26	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 14 August 2001	Examiner Andres, S
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons N : member of the same patent family, corresponding document	

EPO FORM 1533 03 82 (P04C01)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 00 12 6114

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	LIN CHI-TSAI ET AL: "Cloning and characterization of a cDNA for manganese superoxide dismutase from callus of sweet potato." JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY, vol. 45, no. 2, 1997, pages 521-525, XP002174825 ISSN: 0021-8561 * the whole document *	24,25	
X	ZHANG C ET AL: "PHOTOSENSITISATION PROPERTIES OF MITOCHONDRIALLY LOCALISED GREEN FLUORESCENT PROTEIN" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, vol. 242, no. 2, 14 January 1998 (1998-01-14), pages 390-395, XP001002769 ISSN: 0006-291X * the whole document *	24,26	
A	NEUPERT WALTER: "Protein import into mitochondria." ANNUAL REVIEW OF BIOCHEMISTRY, vol. 66, 1997, pages 863-917, XP001015498 ISBN: 0-8243-0866-2 * the whole document *	24-26	
P,X	WO 00 28041 A (MICROBIOLOGICAL RES AUTHORITY ;HALLIS BASSAM (GB); SILMAN NIGEL (G) 18 May 2000 (2000-05-18) * abstract * * page 5, line 10 - line 18 * * page 9, line 28 - page 11, line 6 * * claims; examples 1,2,7,11 *	24-26	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 14 August 2001	Examiner Andres, S
CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document		T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application I: document cited for other reasons S: member of the same patent family, corresponding document	

EPC FORM 1503 03 02 (P40C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 12 6114

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

14-08-2001

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9002804	A	22-03-1990	AU	4195489 A	02-04-1990
WO 0028041	A	18-05-2000	AU	1059400 A	29-05-2000

EPD FORM 20459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)